




# ALGEBRA FOR PREPARATORY TWO FIRST TERM



## Sheet (1)

### Important Revision

[1] Complete by writing the following numbers in the form  $\frac{a}{b}$  where  $a$  &  $b$  are two integers in the simplest form,  $b \neq 0$ :

(1)  $0.2 = \dots\dots\dots$

(2)  $0.3 = \dots\dots\dots$

(3)  $25\% = \dots\dots\dots$

(4)  $|-0.75| = \dots\dots\dots$

(5)  $-6 = \dots\dots\dots$

(6)  $1\frac{1}{4} = \dots\dots\dots$

[2] Complete the following:

(1)  $\sqrt{25 + 144} = \dots\dots\dots$

(2)  $\sqrt{0.25} = \dots\dots\dots$

(3) The standard form of the number  $0.00015$  is  $\dots\dots\dots$

(4) The standard form of the number  $421 \times 10^3$  is  $\dots\dots\dots$

(5) The sum of the two square roots of each number  $2\frac{1}{4} = \dots\dots\dots$

[3] Choose the correct answer:

(1)  $Z^+ \cup \{0\} = \dots\dots\dots$  (  $Z$  ,  $N$  ,  $R$  ,  $Q$  )

(2)  $|-2| + |-4| + |6| = \dots\dots\dots$  (zero ,  $|-12|$  ,  $-12$  ,  $6$  )

(3)  $\sqrt{a^2} = \dots\dots\dots$  (  $a$  ,  $-a$  ,  $|a|$  ,  $\pm a$  )

(4)  $\sqrt{100 - 36} = \dots\dots\dots$  (  $4$  ,  $\pm 4$  ,  $8$  ,  $\pm 8$  )

(5)  $\frac{\sqrt{25-9}}{\sqrt{25}-\sqrt{9}} = \dots\dots\dots$  (  $-1$  ,  $1$  ,  $2$  ,  $3$  )

(6) Which of the following rational numbers lies between  $\frac{1}{5}$  and  $\frac{2}{5}$ ? .....  $(\frac{2}{10}, \frac{1}{10}, 0.3, -0.3)$

(7) The product of the rational number  $\frac{a}{b}$  by its additive inverse equals .....  $(\text{zero}, \frac{-a}{b}, \frac{a^2}{b^2}, \frac{-a^2}{b^2})$

(8)  $3^{10} + 3^{10} + 3^{10} =$  .....  $(3^{10}, 3^{30}, 9^{10}, 3^{11})$

(9) If  $a^{-1} = \frac{2}{3}$ , then  $a =$  .....  $(-\frac{2}{3}, \frac{3}{2}, -\frac{3}{2}, 1)$

(10) The multiplicative inverse of  $5^{-1}$  is .....  $(\frac{1}{5}, 5, -5, \frac{-1}{5})$

[4] Find the value of  $x$  in the following equations:

(1)  $5x + 3 = 20$

.....  
.....

(2)  $5x + 11 = 12$

.....  
.....

(3)  $3x + 5 = 1$

.....  
.....

(4)  $x + 3 = 7$

.....  
.....



[5] Find the solution set of each of the following equations, where  $x \in Q$ :

(1)  $x^2 + 12 = 21$

.....

.....

.....

.....

(2)  $2x^2 - 1 = -9$

.....

.....

.....

.....

(3)  $|x| = 2$

.....

.....

.....

.....

(4)  $\sqrt{x^2} = 4$

.....

.....

.....

.....





## Sheet (2)

### The cube root of a rational number

#### Example (1):

Find the area of a square whose side length 5 cm?

$$\text{Area} = S \times S = S^2 = 5^2 = 25 \text{ cm}^2.$$

The square numbers	The square root
$1^2 = 1$	$\sqrt{1} = 1$
$2^2 = 4$	$\sqrt{4} = 2$
$3^2 = 9$	$\sqrt{9} = 3$
$4^2 = 16$	$\sqrt{16} = 4$
$5^2 = 25$	$\sqrt{25} = 5$
$6^2 = 36$	$\sqrt{36} = 6$
$7^2 = 49$	$\sqrt{49} = 7$
$8^2 = 64$	$\sqrt{64} = 8$
$9^2 = 81$	$\sqrt{81} = 9$
$10^2 = 100$	$\sqrt{100} = 10$
$\sqrt{x^4} = x^2$	$\sqrt{x^6} = x^3$

**Example (2):**

Find the volume of a cube whose edge length 5 cm?

$$V = S \times S \times S = S^3 = 5^3 = 125 \text{ cm}^3.$$

The cub numbers	The cube root
$1^3 = 1$	$\sqrt[3]{1} = 1$
$2^3 = 8$	$\sqrt[3]{8} = 2$
$3^3 = 27$	$\sqrt[3]{27} = 3$
$4^3 = 64$	$\sqrt[3]{64} = 4$
$5^3 = 125$	$\sqrt[3]{125} = 5$
$6^3 = 216$	$\sqrt[3]{216} = 6$
$7^3 = 343$	$\sqrt[3]{343} = 7$
$8^3 = 512$	$\sqrt[3]{512} = 8$
$9^3 = 729$	$\sqrt[3]{729} = 9$
$10^3 = 1000$	$\sqrt[3]{1000} = 10$
$\sqrt[3]{x^3} = x$	$\sqrt[3]{x^6} = x^2$

**[1] Complete the following table:**

Number a	8	125	-27	.....	$3\frac{3}{8}$	$-\frac{8}{125}$	.....	.....
$\sqrt[3]{a}$	.....	.....	.....	-10	.....	.....	6	-4

**[2] Find each of the following:**

(1)  $\sqrt[3]{216} = \dots\dots\dots$

(2)  $\sqrt[3]{-343} = \dots\dots\dots$

(3)  $\sqrt[3]{\frac{64}{125}} = \dots\dots\dots$

(4)  $\sqrt[3]{\frac{-8}{27}} = \dots\dots\dots$

(5)  $\sqrt[3]{0.001} = \dots\dots\dots$

(6)  $\sqrt[3]{-2\frac{10}{27}} = \dots\dots\dots$

(7)  $\sqrt[3]{8x^3} = \dots\dots\dots$

(8)  $\sqrt[3]{-27a^6} = \dots\dots\dots$

**[3] Complete:**

(1)  $\sqrt[3]{x^3} = \dots\dots\dots$

(2)  $\sqrt[3]{\dots\dots\dots} = 4$

(3)  $\sqrt{16} = \sqrt[3]{\dots\dots\dots}$

(4)  $|\sqrt[3]{-125}| = \sqrt{\dots\dots\dots}$

(5)  $\sqrt[3]{8} + \sqrt[3]{-8} = \dots\dots\dots$

(6)  $\sqrt[3]{27} - \sqrt[3]{64} = \dots\dots\dots$

$$(7) \sqrt[3]{27} - \sqrt[3]{-27} = \dots\dots\dots$$

$$(8) \sqrt{9} + \sqrt[3]{-8} = \dots\dots\dots$$

$$(9) \sqrt{64} - \sqrt[3]{64} = \dots\dots\dots$$

$$(10) -\sqrt[3]{-1} - \sqrt{1} = \dots\dots\dots$$

$$(11) \frac{-\sqrt[3]{64}}{\sqrt{64}} = \dots\dots\dots$$

$$(12) \sqrt[3]{64} = \sqrt{\dots\dots\dots}$$

$$(13) \sqrt[3]{64 + \dots\dots\dots} = 5$$

[4] Find the value of  $x$  in each of the following:

$$(1) \sqrt[3]{x} = 5 \quad x = \dots\dots\dots$$

$$(2) \sqrt[3]{x} = \frac{-1}{4} \quad x = \dots\dots\dots$$

$$(3) \sqrt[3]{x} = -\sqrt{4} \quad x = \dots\dots\dots$$

$$(4) \sqrt[3]{x} - 3 = -1 \quad x = \dots\dots\dots$$

$$(5) x^3 = -8 \quad x = \dots\dots\dots$$

$$(6) x^3 = 64 \quad x = \dots\dots\dots$$

$$(7) x^3 + 5 = 32 \quad x = \dots\dots\dots$$

$$(8) 2x^3 = 54 \quad x = \dots\dots\dots$$

$$(9) \frac{1}{5}x^3 = -200 \quad x = \dots\dots\dots$$

**[5] Choose the correct answer from those given:**

- (1)  $\sqrt[3]{(-8)^2} = \dots\dots\dots$  ( 2 , -2 , 4 , -4 )
- (2)  $\sqrt[3]{-64} + \sqrt{16} = \dots\dots\dots$  ( 0 , 8 , -8 ,  $\pm 8$  )
- (3)  $\sqrt{25} - \sqrt[3]{-125} = \dots\dots\dots$  ( 10 , 0 , 5 ,  $\pm 5$  )
- (4)  $\sqrt{(-2)^2} + \sqrt[3]{(-2)^3} = \dots\dots\dots$  ( -4 , 8 , 4 , 0 )
- (5)  $\sqrt[3]{3\frac{3}{8}} + \sqrt{0.25} = \dots\dots\dots$  (  $\frac{3}{2}$  ,  $\frac{1}{2}$  , 2 , -2 )
- (6)  $\sqrt[3]{x} = \frac{1}{4}$ , then  $x = \dots\dots\dots$  (  $\frac{1}{2}$  ,  $\frac{1}{16}$  ,  $\frac{1}{12}$  ,  $\frac{1}{64}$  )
- (7) If the volume of a cube is  $64 \text{ cm}^3$ , then the length of its edge =  $\dots\dots\dots$  cm ( 8 , 4 , 32 , 16 )
- (8) If the capacity of a cubic vessel is 8 litres, then the length of its inner edge is  $\dots\dots\dots$  cm ( 2 , 4 , 20 , 40 )
- (9) If the volume of a sphere is  $36 \pi \text{ cm}^3$ , then the length of its diameter =  $\dots\dots\dots$  cm ( 3 , 6 , 9 , 27 )
- (10) If  $-\sqrt{25} = \sqrt[3]{y}$ , then  $y = \dots\dots\dots$  ( 5 , -5 , 125 , -125 )
- (11) If  $x^3 = 64$ , then  $\sqrt{x} = \dots\dots\dots$  ( 4 , -4 , 2 , -2 )
- (12)  $\sqrt[3]{x^6} = \sqrt{\dots\dots\dots}$  (  $x^3$  ,  $x^2$  ,  $x$  ,  $x^4$  )
- (13) If  $\frac{x}{3} = \frac{9}{x^2}$ , then  $x = \dots\dots\dots$  ( 1 , 3 , 9 , 27 )

[6] Find the S.S. of each of the following equations in Q:

(1)  $x^3 + 27 = 0$

---

---

---

(2)  $8x^3 + 7 = 8$

---

---

---

(3)  $(x + 3)^3 = 343$

---

---

---

(4)  $(5x - 2)^3 + 10 = 18$

---

---

---

(5)  $2x^3 - 5 = x^3 + 3$

---

---

---



**Sheet (3)**  
**The Set of irrational numbers  $Q'$**

The set of irrational numbers denoted by  $Q'$  appear in:

(1) The square root of a non perfect square of a rational number such as:  $\sqrt{2}$ ,  $\sqrt{3}$ ,  $\sqrt{5}$ , .....

(2) The cube root of a non perfect cube of a rational number such as:  $\sqrt[3]{2}$ ,  $\sqrt[3]{3}$ ,  $\sqrt[3]{4}$ , .....

(3)  $\pi \notin Q$

[1] In each of the following, show which of them is a rational number and which of them is an irrational number:

(1) -5

(2)  $2\frac{2}{3}$

(3) 2.06

(4)  $2.3 \times 10^5$

(5)  $-\sqrt{36}$

(6)  $\sqrt[3]{36}$

(7) Zero

(8)  $\sqrt[3]{\frac{-64}{81}}$

(9)  $\sqrt{\frac{1}{3}}$

(10)  $\frac{\pi}{2}$

(11)  $(-5)^{zero}$

(12)  $\sqrt{9} + \sqrt{16}$

(13)  $\sqrt{4} - \sqrt{11}$

(14)  $\sqrt[3]{8} + \sqrt[3]{27}$



[2] Find an approximated value for each of the following numbers:

(1)  $\sqrt{11} \cong \dots\dots\dots$  (to the nearest hundredth)

(2)  $\sqrt[3]{7} \cong \dots\dots\dots$  (to the nearest tenth)

(3)  $\sqrt[3]{-9} \cong \dots\dots\dots$  (to the nearest tenth)

[3] Find two successive integers for each the following numbers to be included between them:

(1)  $\sqrt{5}$  is between  $\dots\dots\dots$  and  $\dots\dots\dots$

(2)  $\sqrt{12}$  is between  $\dots\dots\dots$  and  $\dots\dots\dots$

(3)  $\sqrt[3]{-20}$  is between  $\dots\dots\dots$  and  $\dots\dots\dots$

[4] If  $x$  is an integer, find the value of  $x$  in each of the following cases:

(1)  $x < \sqrt{2} < x + 1$   $x = \dots\dots\dots$

(2)  $x < \sqrt{80} < x + 1$   $x = \dots\dots\dots$

(3)  $x < \sqrt[3]{50} < x + 1$   $x = \dots\dots\dots$

(4)  $x < \sqrt[3]{-100} < x + 1$   $x = \dots\dots\dots$

(5)  $x < |-\sqrt{35}| < x + 1$   $x = \dots\dots\dots$

**[5] Complete the following:**

- (1) The value of  $\sqrt[3]{13}$  to the nearest one decimal is .....
- (2) The two consecutive integers which include the number  $\sqrt{5}$  between them are ..... and .....
- (3)  $\sqrt[3]{x^6} = \sqrt{\dots\dots\dots}$
- (4) The solution set in  $\mathbb{Q}$  of the equation  $5x^2 = 20$  is .....
- (5) If  $x \in \mathbb{Z}$  and  $x < \sqrt[3]{29} < x + 1$ , then  $x = \dots\dots\dots$
- (6) If  $x = \sqrt{3}$ , then  $x^2 = \dots\dots\dots$

**[6] Choose the correct answer:**

- (1) The irrational number in the following numbers is .....  
 (a)  $\sqrt{\frac{1}{4}}$       (b)  $\sqrt[3]{8}$       (c)  $\sqrt{\frac{4}{9}}$       (d)  $\sqrt{2}$
- (2)  $(\sqrt[3]{-3})^3 = \dots\dots\dots$   
 (a) 3      (b) -3      (c)  $\pm 3$       (d)  $\sqrt[3]{-9}$
- (3)  $\sqrt[3]{9} \dots\dots\dots \sqrt{2}$   
 (a) <      (b) >      (c) =      (d)  $\leq$
- (4) The irrational number located between 2 and 3 is .....  
 (a)  $\sqrt{10}$       (b)  $\sqrt{7}$       (c) 2.5      (d)  $\sqrt{3}$

- (5) The irrational number between 3 and 4 is .....  
 (a) 3.6 (b)  $\sqrt{6}$  (c)  $\sqrt{15}$  (d)  $\sqrt{17}$
- (6) The irrational number between -2 and -1 is .....  
 (a) -3 (b)  $-1\frac{1}{2}$  (c)  $-\sqrt{3}$  (d)  $\sqrt{2}$
- (7)  $\sqrt{10} \cong \dots\dots\dots$   
 (a) 2.99 (b) 3.71 (c) 3 (d) -3.2
- (8) The nearest integer to  $\sqrt[3]{26}$  is .....  
 (a) 5 (b) 3 (c) 2 (d) 13
- (9) If  $n \in \mathbb{Z}_+$ ,  $n < \sqrt{26} < n + 1$ , then  $n = \dots\dots\dots$   
 (a) 25 (b) 5 (c) -5 (d) 24
- (10) The area of a square whose side length is  $\sqrt{3}$  cm is .....cm<sup>2</sup>.  
 (a)  $4\sqrt{3}$  (b) 9 (c) 3 (d) 6
- (11) The square whose side length is  $\sqrt{7}$  cm, its area is .....cm<sup>2</sup>.  
 (a) 28 (b) 49 (c) 7 (d) 14
- (12) The square whose area is 10 cm<sup>2</sup>, its side length is ..... cm.  
 (a) 5 (b) -5 (c)  $\sqrt{10}$  (d)  $-\sqrt{10}$
- (13) The S.S. of the equation  $(x - \sqrt{5})(x + \sqrt{3}) = 0$  in  $\mathbb{Q}'$  is .....  
 (a)  $\{\sqrt{5}\}$  (b)  $\{-\sqrt{3}\}$  (c)  $\{-\sqrt{5}, \sqrt{3}\}$  (d)  $\{\sqrt{5}, -\sqrt{3}\}$
- (14)  $|\sqrt[3]{-125}| = \sqrt{\dots\dots\dots}$   
 (a) 25 (b) -25 (c) 5 (d) -5

[7] Find the value of  $x$  in each of the following cases and determine whether  $x \in Q$  or  $x \in Q'$ :

(1)  $5x^2 = 10$

"  $\pm \sqrt{2}$  "

.....

.....

(2)  $4x^2 = 9$

"  $\pm \frac{3}{2}$  "

.....

.....

(3)  $x^3 = 125$

"5"

.....

.....

(4)  $3x^3 = 27$

" $\sqrt[3]{9}$ "

.....

.....

(5)  $0.001x^3 = -8$

" - 20 "

.....

.....

(6)  $(x - 1)^2 = 4$

"3 or - 1"

.....

.....

(7)  $(x - 5)^3 = 1$

"6"

.....

.....



[8] Find in  $\mathbb{Q}$  the S.S. of each of the following equations:

(1)  $x^2 = 13$

.....  
.....  
.....

(2)  $x^3 = 16$

.....  
.....  
.....

(3)  $\frac{2}{5}x^2 = \frac{25}{2}$

.....  
.....  
.....

(4)  $125x^3 - 7 = 20$

.....  
.....  
.....

(5)  $\frac{1}{4}x^2 + 2 = 66$

.....  
.....  
.....



[9] Prove that (V.I):

(1)  $\sqrt{2}$  is included between 1.4 and 1.5

.....  
.....  
.....

(2)  $\sqrt{11}$  is included between 3.31 and 3.32

---

---

---

(3)  $\sqrt{5}$  lies between 2.2 and 2.3

---

---

---

(4)  $\sqrt[3]{15}$  lies between 2.4 and 2.5

---

---

---

(5)  $\sqrt[3]{-17}$  lies between -2.6 and -2.5

---

---

---

(6)  $\sqrt{3} + 1$  lies between 2.7 and 2.8

---

---

---

## Representing an irrational number on the number line

Therefore we can deduce that :

Each irrational number can be represented by a point on the number line.


### Generally

To draw a line segment with length  $\sqrt{a}$  length unit where  $a > 1$ ,

draw a right-angled triangle in which

the length of one side of the right-angle =  $\frac{a-1}{2}$  length unit

and the length of the hypotenuse =  $\frac{a+1}{2}$  length unit.



### Example:

Draw a line segment with length  $= \sqrt{7}$  length unit, then use it to determine the points which represent the following numbers on the number line :


Using the compasses with a distance equal to

the length of  $\overline{BC}$  taking O as a centre

, draw an arc to cut the number line

on the right side of O at the point X,

then X is the point which represents  $\sqrt{7}$



[10] Determine the point that represents each of the following numbers on the number line:

(1)  $\sqrt{3}$

(2)  $-\sqrt{11}$

(3)  $\sqrt{10}$

(4)  $\sqrt{5} + 1$



## Sheet (4)

# The Set of real numbers $\mathbb{R}$

### The set of real numbers

It is the set obtained from the union of the set of rational numbers and the set of irrational numbers. It is denoted by  $\mathbb{R}$


i.e.  $\mathbb{R} = \mathbb{Q} \cup \mathbb{Q}'$  (as shown in the opposite figure)

Noticing that :  $\mathbb{Q} \cap \mathbb{Q}' = \emptyset$

• The opposite Venn diagram shows that :

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$$

$$\text{and } \mathbb{Q}' \subset \mathbb{R}$$



$$\mathbb{R} - \mathbb{Q} = \mathbb{Q}'$$

$$\mathbb{R} - \mathbb{Q}' = \mathbb{Q}$$

$$\mathbb{R}_+ = \{x : x \in \mathbb{R}, x > 0\}$$

$$\mathbb{R}_- = \{x : x \in \mathbb{R}, x < 0\}$$

$$\mathbb{R}_+ \cap \mathbb{R}_- = \emptyset$$

$$\mathbb{R} = \mathbb{R}_+ \cup \{0\} \cup \mathbb{R}_-$$

$$\pi \in \mathbb{Q}'$$

$$\mathbb{R}^* = \mathbb{R} - \{0\} = \mathbb{R}_+ \cup \mathbb{R}_-$$

**[1] Complete the following:**

(1)  $\mathbb{Q} \cap \mathbb{Q}' = \dots\dots\dots$

(2)  $\mathbb{Q} \cup \mathbb{Q}' = \dots\dots\dots$

(3)  $\mathbb{R}_+ \cap \mathbb{R}_- = \dots\dots\dots$

(4)  $\mathbb{R}_+ \cup \mathbb{R}_- = \dots\dots\dots$

(5)  $\mathbb{R} - \mathbb{Q}' = \dots\dots\dots$

(6)  $\mathbb{R} - \mathbb{Q} = \dots\dots\dots$

- (7) The solution set in  $\mathbb{R}$  of the equation  $x^2 + 9 = 0$  is .....
- (8) The cube whose volume is  $8 \text{ cm}^3$ , then the sum of the lengths of its edges is .....cm.
- (9) The two integers which include the number  $\sqrt{12}$  between them are .....and .....
- (10) If  $\sqrt[3]{x} = -5$ , then  $x = \dots\dots\dots$

[2] Put the suitable sign ( $<$ ), ( $>$ ) or ( $=$ ):

- (1)  $\sqrt{5}$   2
- (2)  $\sqrt{7}$   2.6
- (3)  $\sqrt[3]{24}$   3
- (4)  $\sqrt[3]{-24}$   -2
- (5)  $3 - \sqrt{5}$    $\sqrt[3]{-1}$
- (6)  $\sqrt[3]{-8}$    $\sqrt{4}$
- (7)  $1 + \sqrt{3}$    $\sqrt{5}$
- (8)  $\sqrt[3]{3} - 1$   0.2
- (9)  $\sqrt{2} - 1$    $1 - \sqrt{2}$

[3] Choose the correct answer from the given ones:

(1)  $R = \dots\dots\dots$

- (a)  $Q \cup Q'$       (b)  $Z_+ \cup Z_-$       (c)  $R_+ \cup R_-$       (d)  $N \cup R_-$

(2)  $\{x: x \in R, x < 0\} = \dots\dots\dots$

- (a)  $R_+$       (b)  $R_-$       (c)  $R^*$       (d)  $R$

(3) If  $x$  is a negative real number, then which of the following numbers is positive?  $\dots\dots\dots$

- (a)  $x^2$       (b)  $x^3$       (c)  $2x$       (d)  $\frac{x}{2}$

(4)  $R_+ = \dots\dots\dots$

- (a)  $\{x: x \in R, x < 0\}$       (b)  $\{x: x \in R, x \geq 1\}$   
(c)  $\{x: x \in R, x > 0\}$       (d)  $\{x: x \in R, x \geq 0\}$

(5)  $\sqrt[3]{5} \dots\dots \sqrt{3}$

- (a)  $<$       (b)  $>$       (c)  $=$       (d)  $\geq$

(6) The irrational number which is included between 2 and 3 is  $\dots\dots\dots$

- (a)  $\sqrt{10}$       (b)  $\sqrt{7}$       (c) 2.5      (d)  $\sqrt{3}$

(7)  $(-5)^{\text{zero}} = \dots\dots\dots$

- (a) zero      (b) 1      (c) -1      (d) -5

(8) The S.S. of the equation  $x^2 + 1 = 0$  in  $R$  is  $\dots\dots\dots$

- (a)  $\{-1\}$       (b)  $\{1, -1\}$       (c)  $\{1\}$       (d)  $\varphi$

(9)  $\sqrt{(2 - \pi)^2} \dots\dots\dots (2 - \pi)$

- (a)  $<$       (b)  $>$       (c)  $=$       (d)  $\geq$

- (10) If  $x \in R_+$ ,  $y \in R_+$  and  $x^2 > y^2$ , then .....  
 (a)  $x > y$       (b)  $x < y$       (c)  $x = y$       (d)  $x \leq y$

- (11) If  $\frac{1}{a}$  and  $\frac{a}{\sqrt{5}}$  are real numbers included between 0 and 1,  
 then  $a =$  .....  
 (a) -2      (b) 1      (c)  $\sqrt{5}$       (d) 2

**[4] Arrange the following numbers in an ascending order:**

- (1)  $\sqrt{8}$ ,  $-\sqrt{3}$ ,  $\sqrt{15}$ ,  $\sqrt{5}$ ,  $-\sqrt{7}$  and  $-\sqrt{11}$

The order is:

.....

- (2)  $\sqrt{27}$ ,  $-\sqrt{45}$ ,  $\sqrt{20}$ , 0.6 and  $\sqrt[3]{-1}$

The order is:

.....

**[5] Arrange the following numbers in a descending order:**

- (1)  $\sqrt{62}$ , 8,  $-\sqrt{50}$  and  $\sqrt{70}$

The order is:

.....

- (2)  $\sqrt{6}$ , 9,  $-\sqrt{10}$ ,  $-\sqrt{7}$ ,  $-\sqrt{50}$  and  $\sqrt{101}$

The order is:

.....

**[6] Write three positive irrational numbers less than 2:**

.....

[7] Write three negative irrational numbers greater than  $-\sqrt{6}$ :

.....



[8] Write four irrational numbers included between 15 and 17:

.....



[9] Write three irrational positive numbers less than 3:

.....



[10] Solve the following equations to the nearest hundredth given  $x \in R$ :

(1)  $x^2 - 6 = 0$

.....  
 .....  
 .....

(2)  $\frac{3}{4}x^2 = 24$

.....  
 .....  
 .....

(3)  $\frac{1}{2}x^2 - 5 = 0$

.....  
 .....  
 .....

(4)  $5x^3 + 3 = 2$

.....  
 .....  
 .....

$$(5) (x^2 - 9)(x^3 - 5) = 0$$

.....

.....

.....

$$(6) (2x^3 - 5)(x^2 + 1) = 0$$

.....

.....

.....

### Rules to solve geometric applications

#### The cube

Let the edge length =  $l$  ,

volume =  $l^3$  , lateral area =  $4l^2$  , total area =  $6l^2$

[11] Find the edge length of a cube whose volume is  $1.728 \text{ cm}^3$ . Is the edge length a rational number? " $\frac{6}{5} \text{ cm}$ "

.....

.....

.....

[12] A cube whose total area is  $13.5 \text{ cm}^2$ . Find its edge length. Is the edge length a rational number? " $1.5 \text{ cm}$ "

.....

.....

.....

## The Square

Let the edge length =  $S$ , then

$$\text{area} = S^2$$

$$S = \sqrt{\text{area}}$$

Let the diagonal length =  $d$ , then  $\text{area} = \frac{1}{2} d^2$ ,  
 $d = \sqrt{2 \times \text{area}}$  or  $d = \sqrt{2 \times s \times s}$

[13] Find the side length of a square whose area is 5 cm<sup>2</sup>. Is the edge length a rational number? " $\sqrt{5}\text{cm}$ "

.....

.....

.....

[14] A square is of area 32 cm<sup>2</sup>, Find its side length and its diagonal length? " $\sqrt{32}, 8$ "

.....

.....

.....

[15] A square is of side length 6 cm. Find its diagonal length? " $\sqrt{72}\text{cm}$ "

.....

.....

.....





## Sheet (5) Subsets of real numbers (Intervals)

If  $X$  is the set of integers which greater than or equal to  $-3$  and less than  $2$ , we can express the set  $X$  by the description method as follows:

$$X = \{a: a \in \mathbb{Z}, -3 \leq a < 2\}$$

we can express it by listing method









$$X = \{-3, -2, -1, 0, 1, 2\}$$

If  $K$  is the set of real number which greater than or equal to  $-3$  and less than  $2$ , we can express the set  $K$  by the description method as follows:

$$K = \{a: a \in \mathbb{R}, -3 \leq a < 2\}$$

but it is impossible to express the set  $K$  by listing method because there are an infinity of real numbers between  $-3$  and  $2$ . So we use another method to express a subset of the set of real numbers, which is (the intervals).

### Types of intervals

Types of intervals		The interval	Expression by distinguished property	Representation on the number line	Notice that
The limited intervals	Closed	$[a, b]$	$\{x : x \in \mathbb{R}, a \leq x \leq b\}$		<ul style="list-style-type: none"><li><math>a \in [a, b]</math></li><li><math>b \in [a, b]</math></li></ul>
	Opened	$]a, b[$	$\{x : x \in \mathbb{R}, a < x < b\}$		<ul style="list-style-type: none"><li><math>a \notin ]a, b[</math></li><li><math>b \notin ]a, b[</math></li></ul>
	half opened (half closed)	$[a, b[$	$\{x : x \in \mathbb{R}, a \leq x < b\}$		<ul style="list-style-type: none"><li><math>a \in [a, b[</math></li><li><math>b \notin [a, b[</math></li></ul>
		$]a, b]$	$\{x : x \in \mathbb{R}, a < x \leq b\}$		<ul style="list-style-type: none"><li><math>a \notin ]a, b]</math></li><li><math>b \in ]a, b]</math></li></ul>
The unlimited intervals		$[a, \infty[$	$\{x : x \in \mathbb{R}, x \geq a\}$		$a \in [a, \infty[$
		$]a, \infty[$	$\{x : x \in \mathbb{R}, x > a\}$		$a \notin ]a, \infty[$
		$]-\infty, a]$	$\{x : x \in \mathbb{R}, x \leq a\}$		$a \in ]-\infty, a]$
		$]-\infty, a[$	$\{x : x \in \mathbb{R}, x < a\}$		$a \notin ]-\infty, a[$

Remarks

1  $\mathbb{R} = ]-\infty, \infty[$









2  $\mathbb{R}_+ = ]0, \infty[$

3  $\mathbb{R}_- = ]-\infty, 0[$

4 The set of non-negative real numbers  $= \mathbb{R}_+ \cup \{0\} = [0, \infty[$

5 The set of non-positive real numbers  $= \mathbb{R}_- \cup \{0\} = ]-\infty, 0]$

[1] Complete the following table:

(1)	$[-1, 2]$	$\{x: x \in \mathbb{R}, -1 \leq x \leq 2\}$	
(2)	$[1, 3[$	.....	
(3)	.....	$\{x: x \in \mathbb{R}, 0 < x \leq 3\}$	
(4)	.....	.....	
(5)	$] -\infty, 1]$	.....	
(6)	.....	.....	
(7)	.....	$\{x: x \in \mathbb{R}, x < 4\}$	
(8)	$[-2, \infty[$	.....	

[2] Complete using  $(\in)$  or  $(\notin)$ :

(1) 3 .....  $[3, 5]$  (2) -2 .....  $] -2, 1]$

(3) 0 .....  $[-1, 4[$  (4)  $|-3|$  .....  $[2, \infty[$

(5)  $\sqrt{9}$  .....  $] -3, \infty[$  (6)  $\sqrt[3]{-1}$  .....  $] -\infty, 1]$

(7)  $1.3 \times 10^{-5}$  .....  $\mathbb{R}_+$  (8)  $\sqrt{2}$  .....  $[2, 5]$

(9) 5 .....  $] \sqrt{5}, \sqrt{23} [$  (10)  $\sqrt[3]{-125}$  .....  $] -\sqrt{25}, \sqrt{25}]$

**[3] Choose the correct answer:**

<b>1</b>	$\sqrt[3]{8} \dots\dots\dots ]-\infty, 4[$ (a) $\in$ (b) $\notin$ (c) $\subset$ (d) $\not\subset$
<b>2</b>	$5 \in \dots\dots\dots$ (a) $]5, \infty[$ (b) $] -\infty, 5[$ (c) $(3, 5)$ (d) $[-5, \infty[$
<b>3</b>	The opposite figure represents the interval $\dots\dots\dots$ (a) $[-4, 8[$ (b) $[8, -4]$ (c) $[-4, 8]$ (d) $]-4, 8[$
<b>4</b>	$\mathbb{R} = \dots\dots\dots$ (a) $\mathbb{R}_+ \cap \mathbb{R}_-$ (b) $\mathbb{R}_+ \cup \mathbb{R}_-$ (c) $] -\infty, \infty[$ (d) $\mathbb{Q} \cap \mathbb{Q}$
<b>5</b>	$\mathbb{R}_+ = \dots\dots\dots$ (a) $]0, \infty[$ (b) $] -\infty, 0[$ (c) $[0, \infty[$ (d) $] -\infty, 0]$
<b>6</b>	$\mathbb{R}_- = \dots\dots\dots$ (a) $]0, \infty[$ (b) $] -\infty, 0[$ (c) $[0, \infty[$ (d) $] -\infty, 0]$
<b>7</b>	The set of non-negative real numbers = $\dots\dots\dots$ (a) $]0, \infty[$ (b) $] -\infty, 0[$ (c) $[0, \infty[$ (d) $] -\infty, 0]$
<b>8</b>	The set of non-positive real numbers = $\dots\dots\dots$ (a) $]0, \infty[$ (b) $] -\infty, 0[$ (c) $[0, \infty[$ (d) $] -\infty, 0]$



# Operations on Intervals

[1] Complete:

1  $[1, 3] \cup [2, 5[ = \dots\dots\dots$



2  $]1, 3] \cup [2, 5] = \dots\dots\dots$



3  $]-\infty, 1] \cup [-4, \infty[ = \dots\dots\dots$



4  $]3, 5[ \cup \{3, 5\} = \dots\dots\dots$

5  $]-2, 2] \cup \{-2, 0\} = \dots\dots\dots$

6  $]5, 7[ \cup \{5, 7\} = \dots\dots\dots$

7  $\mathbb{N} \cap ]1, 2[ = \dots\dots\dots$

8  $]1, 7[ \cap ]3, 5[ = \dots\dots\dots$



9  $[-3, 1[ \cap [-1, 4[ = \dots\dots\dots$



10  $[-2, 5] \cap ]4, 6] = \dots\dots\dots$



11  $] -3, 5] \cap [0, 3[ = \dots\dots\dots$



12  $[1, 5] - \{1, 5\} = \dots\dots\dots$

13  $[2, 5] - \{5\} = \dots\dots\dots$

14  $[2, 5] - \{2, 5\} = \dots\dots\dots$

15  $[3, 4] - \{3, 4\} = \dots\dots\dots$

16  $[3, 5] - \{3\} = \dots\dots\dots$

17  $[3, 7[ - ] -2, 5] = \dots\dots\dots$



18  $[-4, 6] - ] -4, 6[ = \dots\dots\dots$



19  $[-1, 5] - ] -1, 5[ = \dots\dots\dots$



20  $[2, 7] - ]2, 7[ = \dots\dots\dots$





**[2] Essay problems:**

- 1** If  $X = [-1, 4]$ ,  $Y = [3, \infty[$ ,  $Z = \{3, 4\}$ , find using the number line :
- (1)  $X \cup Y$                       (2)  $X \cap Y$                       (3)  $X - Z$



.....

.....

.....

- 2** If  $X = [3, \infty[$ ,  $Y = ]-4, 8[$   
Find : (1)  $X \cup Y$                       (2)  $X \cap Y$                       (3)  $\bar{X}$



.....

.....

.....

- 3** Find each of the following :
- (1)  $[0, 5] \cup [3, 8[$                       (2)  $[1, 5] \cap ]-2, 3]$



.....

.....

- 4** If  $X = [-2, 3]$ ,  $Y = [1, 5[$ , then find by using the number line :  $X \cup Y$ ,  $X - Y$



.....

.....

**[1] Choose the correct answer:**

<b>1</b>	$] -1, 3] \cap [-3, -1] = \dots\dots\dots$ (a) $\emptyset$ (b) $\{-3\}$ (c) $\{-1\}$ (d) $\{3\}$
<b>2</b>	$[1, 5] \cap ] -2, 3] = \dots\dots\dots$ (a) $\{1, 3\}$ (b) $]1, 3[$ (c) $[1, 3]$ (d) $[1, 3[$
<b>3</b>	$] -3, 5[ \cap [0, 3[ = \dots\dots\dots$ (a) $[0, 3]$ (b) $[0, 3[$ (c) $] -3, 0[$ (d) $[3, 5[$
<b>4</b>	$[2, 7] - \{2, 7\} = \dots\dots\dots$ (a) $[1, 6]$ (b) $\emptyset$ (c) $]2, 7[$ (d) $\{0\}$
<b>5</b>	$[-2, 5] - \{-2, 6\} = \dots\dots\dots$ (a) $] -2, 5[$ (b) $] -2, 6[$ (c) $] -2, 5]$ (d) $[-2, 5[$
<b>6</b>	$[-3, 7] - \{-3, 7\} = \dots\dots\dots$ (a) $[-3, 7[$ (b) $] -3, 7]$ (c) $] -3, 7[$ (d) $(0, 0)$

**[2] Essay problems:**

- 1** If  $X = [-1, 4]$ ,  $Y = [3, \infty[$ ,  $Z = \{3, 4\}$ , find using the number line :
- (1)  $X \cup Y$  (2)  $X \cap Y$  (3)  $X - Z$



.....

.....

.....

- 2** If  $X = [-2, 1]$  and  $Y = [0, \infty[$

**Find :** (1)  $X \cap Y$  (2)  $X \cup Y$



.....

.....



**3** If  $X = [3, \infty[$ ,  $Y = ]-4, 8[$

Find : (1)  $X \cup Y$

(2)  $X \cap Y$

(3)  $\bar{X}$



**4** If  $X = [-1, 4]$  and  $Y = [2, 7]$ , then find each of :

(1)  $X \cap Y$

(2)  $Y \cup X$



**5** If  $X = [-2, 1]$ ,  $Y = [0, \infty[$

Find : (1)  $X \cap Y$

(2)  $X \cup Y$

(3)  $Y - X$



**6** If  $X = [-1, 4]$ ,  $Y = [3, \infty[$ , find using the number line each of :

(1)  $X \cup Y$


(2)  $X - Y$



**7** Find each of the following :

(1)  $[0, 5] \cup [3, 8[$

(2)  $[1, 5] \cap ]-2, 3]$



**8** If  $X = [-2, 3]$ ,  $Y = [1, 5[$ , then find by using the number line :  $X \cup Y$ ,  $X - Y$



**9** If  $X = [-2, 4]$  and  $Y = ]2, \infty[$ , find each of the following using the number line :

(1)  $X \cap Y$

(2)  $X - Y$



**10** If  $X = [-1, 4]$ ,  $Y = [2, 7]$ , then find each of the following by using the number line :

(1)  $X \cup Y$

(2)  $X \cap Y$

(3)  $X - Y$



**11** If  $X = [1, 5]$ ,  $Y = [2, 7]$  Find by using the number line :

(1)  $X \cap Y$

(2)  $X \cup Y$



## Sheet (6)

# Operations on the real numbers

We know that  $2x$  and  $3x$  are two like algebraic terms, then  $2x + 3x = 5x$

then we deduce that  $2\sqrt{5} + 3\sqrt{5} = (2 + 3)\sqrt{5} = 5\sqrt{5}$ ,

but  $2x$  and  $3y$  are two unlike algebraic terms then their sum  $2x + 3y$

then the sum of  $2\sqrt{3}$ ,  $3\sqrt{2}$  written in the form  $2\sqrt{3} + 3\sqrt{2}$

### Properties of addition of real numbers

#### Closure :

For every  $a \in \mathbb{R}$  and  $b \in \mathbb{R}$  we find that  $(a + b) \in \mathbb{R}$

**i.e.** The sum of any two real numbers is a real number, therefore we say  $\mathbb{R}$  is closed under addition operation.

For example :

•  $\sqrt{5} \in \mathbb{R}$  and  $2\sqrt{5} \in \mathbb{R}$  we find that :  $\sqrt{5} + 2\sqrt{5} = 3\sqrt{5} \in \mathbb{R}$

#### Commutative property :

For every  $a \in \mathbb{R}$  and  $b \in \mathbb{R}$  it will be  $a + b = b + a$

For example :

$$5\sqrt[3]{2} + 4\sqrt[3]{2} = 9\sqrt[3]{2} \text{ , } 4\sqrt[3]{2} + 5\sqrt[3]{2} = 9\sqrt[3]{2}$$

**i.e.**  $5\sqrt[3]{2} + 4\sqrt[3]{2} = 4\sqrt[3]{2} + 5\sqrt[3]{2}$

#### Associative property :

For every  $a \in \mathbb{R}$  ,  $b \in \mathbb{R}$  and  $c \in \mathbb{R}$  it will be  $(a + b) + c = a + (b + c) = a + b + c$

For example :

$$(\sqrt{3} + 2\sqrt{3}) + 5\sqrt{3} = 3\sqrt{3} + 5\sqrt{3} = 8\sqrt{3} \text{ ,}$$

$$\sqrt{3} + (2\sqrt{3} + 5\sqrt{3}) = \sqrt{3} + 7\sqrt{3} = 8\sqrt{3}$$

**i.e.**  $(\sqrt{3} + 2\sqrt{3}) + 5\sqrt{3} = \sqrt{3} + (2\sqrt{3} + 5\sqrt{3})$

**The additive neutral :**

For every  $a \in \mathbb{R}$  it will be  $a + 0 = 0 + a = a$

**i.e.** Zero is the additive neutral.

For example :  $\sqrt{2} + 0 = 0 + \sqrt{2} = \sqrt{2}$  ,  $-\sqrt[3]{5} + 0 = 0 + (-\sqrt[3]{5}) = -\sqrt[3]{5}$

**The additive inverse of every real number :**

For every  $a \in \mathbb{R}$  there is  $(-a) \in \mathbb{R}$  where  $a + (-a) = \text{zero (the additive neutral)}$

For example :

- The additive inverse of the number  $\sqrt{3}$  is  $-\sqrt{3}$  and vice versa because  $\sqrt{3} + (-\sqrt{3}) = 0$
- The additive inverse of the number  $2 + \sqrt{5}$  is  $-(2 + \sqrt{5})$  and equals  $-2 - \sqrt{5}$
- The additive inverse of the number  $3 - \sqrt{2}$  is  $-(3 - \sqrt{2})$  and equals  $\sqrt{2} - 3$
- The additive inverse of the number zero is itself.

**[1] Find the result of each of the following in the simplest form:**

(1)  $\sqrt{3} + 2\sqrt{3} = \dots\dots\dots$

(2)  $3\sqrt{2} - 5\sqrt{2} = \dots\dots\dots$

(3)  $2\sqrt{5} - 3\sqrt{5} + \sqrt{5} = \dots\dots\dots$

(4)  $5\sqrt[3]{7} - 8\sqrt[3]{7} + 2\sqrt[3]{7} = \dots\dots\dots$

(5)  $4\sqrt{5} - 2\sqrt{5} + 5\sqrt{5} - \sqrt{5} = \dots\dots\dots$

(6)  $5\sqrt{3} - 7\sqrt{3} + 3\sqrt{3} - \sqrt{3} = \dots\dots\dots$

(7)  $\sqrt{5} - \sqrt{3} + 2\sqrt{5} + \sqrt{3} = \dots\dots\dots$

(8)  $2\sqrt{3} + 5 + \sqrt{3} - 6 = \dots\dots\dots$

$$(9) \quad 2\sqrt{7} - 3\sqrt{2} + \sqrt{7} + 5\sqrt{7} = \dots\dots\dots$$

$$(10) \quad 2\sqrt{2} - 3\sqrt[3]{2} + 5\sqrt{2} + \sqrt[3]{2} = \dots\dots\dots$$

$$(11) \quad 8\sqrt{\frac{1}{4}} + 2\sqrt[3]{3} - \sqrt[3]{64} - 5\sqrt[3]{3} = \dots\dots\dots$$

$$(12) \quad \frac{1}{4}\sqrt{2} + \frac{2}{7}\sqrt{5} + \frac{3}{4}\sqrt{2} - \frac{2}{7}\sqrt{5} = \dots\dots\dots$$

### The properties of multiplication operation of real numbers

#### Closure :

For every  $a \in \mathbb{R}$  and  $b \in \mathbb{R}$  it will be  $a \times b \in \mathbb{R}$

**i.e.** The product of any two real numbers is a real number therefore we say :

The multiplication operation is closed in  $\mathbb{R}$

For example :

$$\bullet \sqrt{3} \in \mathbb{R} \text{ and } 2\sqrt{3} \in \mathbb{R}$$

$$\text{We find that : } \sqrt{3} \times 2\sqrt{3} = 2 \times 3 = 6 \in \mathbb{R}$$

#### Commutative property :

For every  $a \in \mathbb{R}$  and  $b \in \mathbb{R}$  it will be  $a \times b = b \times a$

For example :

$$\bullet 2\sqrt{5} \times 3\sqrt{5} = 6 \times 5 = 30, \quad 3\sqrt{5} \times 2\sqrt{5} = 6 \times 5 = 30$$

$$\text{i.e. } 2\sqrt{5} \times 3\sqrt{5} = 3\sqrt{5} \times 2\sqrt{5}$$

#### The associative property :

For every  $a \in \mathbb{R}$ ,  $b \in \mathbb{R}$  and  $c \in \mathbb{R}$  it will be  $(a \times b) \times c = a \times (b \times c) = a \times b \times c$

For example :

$$\bullet (2\sqrt{7} \times 4\sqrt{7}) \times \sqrt{7} = 56 \times \sqrt{7} = 56\sqrt{7}$$

$$\bullet 2\sqrt{7} \times (4\sqrt{7} \times \sqrt{7}) = 2\sqrt{7} \times 28 = 56\sqrt{7}$$

$$\text{i.e. } (2\sqrt{7} \times 4\sqrt{7}) \times \sqrt{7} = 2\sqrt{7} \times (4\sqrt{7} \times \sqrt{7})$$



**The multiplicative neutral :**

For every  $a \in \mathbb{R}$  it will be  $a \times 1 = 1 \times a = a$

**i.e.** One is the multiplicative neutral in  $\mathbb{R}$

For example :

•  $\sqrt[3]{5} \times 1 = 1 \times \sqrt[3]{5} = \sqrt[3]{5}$

**The multiplicative inverse of any non-zero real number :**

For every real number  $a \neq 0$  , there is a real number  $\frac{1}{a}$  where  $a \times \frac{1}{a} = 1$  which is the multiplicative neutral.

For example :

- The multiplicative inverse of  $\sqrt{3}$  is  $\frac{1}{\sqrt{3}}$   
because  $\sqrt{3} \times \frac{1}{\sqrt{3}} = 1$
- The multiplicative inverse of  $-\frac{\sqrt{2}}{5}$  is  $-\frac{5}{\sqrt{2}}$
- The multiplicative inverse of the number 1 is itself and also the multiplicative inverse of  $-1$  is itself.

**Notice that :**

Both the number and its multiplicative inverse have the same sign.

**Notice that :**

There is no multiplicative inverse for the number zero because  $\frac{1}{\text{zero}}$  is meaningless (**i.e.** undefined)

**Remark**

- Since each non-zero real number has a multiplicative inverse then the division operation by any real number does not equal zero is possible in  $\mathbb{R}$  and it is defined as

For every  $a \in \mathbb{R}$  and  $b \in \mathbb{R}^*$  it will be  $a \div b = a \times \frac{1}{b}$

**i.e.** The division operation  $(a \div b)$  means multiplying the number  $a$  by the multiplicative inverse of the number  $b$  such that  $b \neq 0$

**Then we can deduce that :**

The division operation in  $\mathbb{R}$  is not commutative and it is not associative.

[2] Find the result of each of the following in the simplest form:

$$(1) \quad \sqrt{3} \times \sqrt{3} = \dots\dots\dots$$

$$(2) \quad -2\sqrt{5} \times 3\sqrt{5} = \dots\dots\dots$$

$$(3) \quad 2 \times 3\sqrt{2} = \dots\dots\dots$$

$$(4) \quad \frac{1}{3} \sqrt{3} \times \sqrt{3} = \dots\dots\dots$$

$$(5) \quad (\sqrt[3]{5})^3 \times 3\sqrt{3} = \dots\dots\dots$$

$$(6) \quad 2\sqrt{3} \times \frac{2\sqrt{7}}{7} \div \frac{20\sqrt{3}}{5\sqrt{7}} = \dots\dots\dots$$

[3] Make the denominator in each of the following an integer:

$$(1) \quad \frac{3}{\sqrt{3}} = \dots\dots\dots$$

$$(2) \quad \frac{10}{\sqrt{5}} = \dots\dots\dots$$

$$(3) \quad \frac{-6}{\sqrt{3}} = \dots\dots\dots$$

$$(4) \quad \frac{6}{2\sqrt{3}} = \dots\dots\dots$$

$$(5) \quad \frac{\sqrt{2} + 3}{\sqrt{2}} = \dots\dots\dots$$



## Distributing multiplication on addition and subtraction

For any three real numbers  $a$ ,  $b$  and  $c$  it will be :

$$\bullet a(b \pm c) = ab \pm ac$$

$$\bullet (b \pm c)a = ba \pm ca$$

### Remarks:

- $(a + b)(a - b) = a^2 - b^2$
- $(a + b)^2 = a^2 + 2ab + b^2$
- $(a - b)^2 = a^2 - 2ab + b^2$

[4] Find the result of each of the following in the simplest form:

(1)	$2(\sqrt{2} + \sqrt{5}) =$ .....
(2)	$\sqrt{2}(5 + \sqrt{2}) =$ .....
(3)	$\sqrt{7}(\sqrt{7} + 2) =$ .....
(4)	$-\sqrt{3}(-5 - \sqrt{3})$ .....
(5)	$-2\sqrt{5}(3 - \sqrt{5}) =$ .....
(6)	$\sqrt{7}\left(\frac{2}{\sqrt{7}} - \sqrt{7} + 3\right) =$ .....

[5] Find the result of each of the following operations:

(1)	$(\sqrt{2} + 1)(\sqrt{2} - 1) =$ .....
(2)	$(4 - 3\sqrt{2})(4 + 3\sqrt{2}) =$ .....
(3)	$(\sqrt{5} - 1)^2 =$ .....
(4)	$(2\sqrt{3} + 4)^2 =$ .....

**[6] Complete the following:**

(1)	The multiplicative neutral in $\mathbb{R}$ is ..... and the additive neutral in $\mathbb{R}$ is .....
(2)	The additive inverse of the number $1 - \sqrt{2}$ is .....
(3)	The multiplicative inverse of the number $\frac{2\sqrt{3}}{5}$ is $\frac{\dots\dots\dots}{6}$
(4)	The multiplicative inverse of the number $\frac{3}{\sqrt{3}}$ is $\frac{\dots\dots\dots}{\sqrt{3}}$
(5)	If : $a = \sqrt{5}$ and $b = 2\sqrt{5}$ , then : $ab = \dots\dots\dots$
(6)	If : $x = \sqrt{5} + 2$ and $y = \sqrt{5} - 2$ then $(x + y)^2 = \dots\dots\dots$
(7)	If : $x = 2\sqrt[3]{5}$ , then $x^3 = \dots\dots\dots$
(8)	The solution set of the equation : $x^2 + 25 = 0$ in $\mathbb{R}$ is .....
(9)	$\mathbb{R}_+ \cup [-3, 2[ = \dots\dots\dots$

**[7] Choose the correct answer:**

(1)	$2\sqrt{5} + 3\sqrt{5} = \dots\dots\dots$ (a) $5\sqrt{10}$ (b) $5\sqrt{5}$ (c) $6\sqrt{5}$ (d) 30
(2)	The multiplicative inverse of the number $\frac{\sqrt{3}}{6}$ is ..... (a) $\frac{\sqrt{6}}{3}$ (b) $2\sqrt{3}$ (c) $\frac{3}{\sqrt{6}}$ (d) $-\frac{\sqrt{3}}{6}$
(3)	$\sqrt{3} + (-\sqrt{3}) = \dots\dots\dots$ (a) $2\sqrt{3}$ (b) $2\sqrt{6}$ (c) $\sqrt{6}$ (d) zero
(4)	$\square - 2\sqrt{3} \times \sqrt{3} = \dots\dots\dots$ (a) - 6                      (b) $-2\sqrt{3}$ (c) $2\sqrt{3}$ (d) 6

(5)	The additive inverse of the number $\frac{6}{\sqrt{2}} = \dots\dots\dots$ (a) $-2\sqrt{3}$ (b) $2\sqrt{3}$ (c) $-3\sqrt{2}$ (d) $3\sqrt{2}$
(6)	The additive inverse of the number $(\sqrt{2} - \sqrt{5}) = \dots\dots\dots$ (a) $\sqrt{2} + \sqrt{5}$ (b) $\sqrt{5} - \sqrt{2}$ (c) $\sqrt{2} - \sqrt{5}$ (d) $-\sqrt{2} - \sqrt{5}$
(7)	The multiplicative inverse of the number $\sqrt{5}$ is $\dots\dots\dots$ (a) $-5$ (b) $\frac{-1}{5}$ (c) $\frac{5}{\sqrt{5}}$ (d) $\frac{\sqrt{5}}{5}$
(8)	$(\sqrt{5} + 3\sqrt{5}) \div \sqrt{5} = \dots\dots\dots$ (a) $3\sqrt{5}$ (b) $3$ (c) $5$ (d) $4$
(9)	If : $x = \sqrt{2} + 10$ , $y = \sqrt{2} - 10$ , then $(x + y)^2 = \dots\dots\dots$ (a) $4$ (b) $6$ (c) $8$ (d) $4\sqrt{2}$
(10)	$[2, 5] - \{2, 5\} = \dots\dots\dots$ (a) $[3, 4]$ (b) $]2, 5[$ (c) $\{2, 5\}$ (d) $[2, 5]$
(11)	If : $x^3 + 9 = 1$ where $x \in \mathbb{R}$ , then $x = \dots\dots\dots$ (a) $-8$ (b) $-2$ (c) $2$ (d) $8$
(12)	If : $x = \sqrt{3} + 2$ , then $x^2 = \dots\dots\dots$ (a) $5$ (b) $7$ (c) $7 + 2\sqrt{3}$ (d) $7 + 4\sqrt{3}$
(13)	If : $x^2 - y^2 = 60$ , $x + y = 5\sqrt{6}$ , then $x - y = \dots\dots\dots$ (a) $\sqrt{6}$ (b) $2\sqrt{6}$ (c) $3\sqrt{6}$ (d) $4\sqrt{6}$

[8] If  $x = \sqrt{5} - 2$  and  $y = \sqrt{5} + 2$ , find the value of:

(1)  $x + y = \dots\dots\dots$

(2)  $x - y = \dots\dots\dots$

(3)  $xy = \dots\dots\dots$

[9] If  $x = -\sqrt{3}$  and  $y = 2\sqrt{3} - 3$ , find the value of:

(1)  $x + y =$  .....

(2)  $xy =$  .....

(3)  $\frac{y}{x} =$  .....



## Sheet (7)

# Operations on the square roots

If a and b are two non negative real numbers , then

$$\boxed{1 \quad \sqrt{a} \times \sqrt{b} = \sqrt{ab}}$$

For example :

$$\bullet \sqrt{3} \times \sqrt{12} = \sqrt{36} = 6$$

$$\bullet \sqrt{50} = \sqrt{25 \times 2} = \sqrt{25} \times \sqrt{2} = 5\sqrt{2}$$

$$\boxed{2 \quad \frac{\sqrt{a}}{\sqrt{b}} = \sqrt{\frac{a}{b}} \text{ where } b \neq 0}$$

For example :

$$\bullet \frac{\sqrt{8}}{\sqrt{2}} = \sqrt{\frac{8}{2}} = \sqrt{4} = 2$$

$$\bullet \sqrt{\frac{16}{49}} = \frac{\sqrt{16}}{\sqrt{49}} = \frac{4}{7}$$

$$\boxed{3 \quad \frac{\sqrt{a}}{\sqrt{b}} = \frac{\sqrt{a}}{\sqrt{b}} \times \frac{\sqrt{b}}{\sqrt{b}} = \frac{\sqrt{ab}}{b} \text{ where } b \neq 0}$$

This operation is carried out to make the denominator an integer.

For example :

$$\bullet \frac{\sqrt{2}}{\sqrt{5}} = \frac{\sqrt{2}}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}} = \frac{\sqrt{10}}{5}$$

$$\bullet \sqrt{\frac{3}{2}} = \frac{\sqrt{3}}{\sqrt{2}} = \frac{\sqrt{3}}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{\sqrt{6}}{2}$$

[1] Put each of the following in the form  $a\sqrt{b}$  where a and b are two integers and b is the least possible value:

(1)  $\sqrt{12} = \sqrt{4 \times 3} = 2\sqrt{3}$

(2)  $\sqrt{28} = \sqrt{4 \times 7} = 2\sqrt{7}$

(3)  $2\sqrt{72} = 2 \times \sqrt{36 \times 2} = 2 \times 6\sqrt{2} = 12\sqrt{2}$

(4)  $\frac{2}{5}\sqrt{1000} = \frac{2}{5}\sqrt{100 \times 10} = \frac{2}{5} \times 10\sqrt{10} = 4\sqrt{10}$

$$(5) 2\sqrt{\frac{1}{2}} = \sqrt{2^2 \times \frac{1}{2}} = \sqrt{2} \left( x\sqrt{\frac{1}{x}} = \sqrt{x} \right)$$

$$(6) 6\sqrt{\frac{2}{3}} = \sqrt{36 \times \frac{2}{3}} = 2\sqrt{6}$$

[2] Simplify each of the following to the simplest form:

$$(1) \sqrt{50} + \sqrt{8}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(2) 3\sqrt{2} + \sqrt{8} - \sqrt{18}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(3) \sqrt{98} - \sqrt{128} - \sqrt{18} + 4\sqrt{2}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(4) \sqrt{27} + 5\sqrt{18} - \sqrt{300}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(5) 2\sqrt{18} + \sqrt{50} + \frac{1}{3}\sqrt{162}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(6) 2\sqrt{5} + 4\sqrt{20} - 5\sqrt{\frac{1}{5}}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(7) 2\sqrt{5} + 6\sqrt{\frac{1}{3}} - \sqrt{12} - 5\sqrt{\frac{1}{5}}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(8) \sqrt{18} - \frac{\sqrt{12}}{\sqrt{6}}$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

(9)  $2\sqrt{3} \times 5\sqrt{2}$

= .....

= .....

(10)  $\sqrt{5} \times 2\sqrt{10}$

= .....

= .....


(11)  $\sqrt{\frac{2}{7}} \times \sqrt{\frac{7}{2}} =$  .....

(12)  $\frac{3\sqrt{15}}{\sqrt{5}} =$  .....

(13)  $12\sqrt{\frac{2}{3}} \times \sqrt{54}$

= .....

= .....



[3] Simplify each of the following to the simplest form:

(1)  $\sqrt{6}(\sqrt{3} - \sqrt{2}) =$  .....

(2)  $(3\sqrt{5} - \sqrt{7})(3\sqrt{5} + \sqrt{7}) =$  .....

(3)  $(3\sqrt{2} - 5)(3\sqrt{2} + 5) =$  .....

(4)  $(\sqrt{2} + \sqrt{6})^2 =$  .....

(5)  $(\sqrt{3} - \sqrt{2})^2 =$  .....



**[4] Choose the correct answer:**

(1)	$\frac{\sqrt{63}}{\sqrt{7}} = \dots\dots\dots$	(a) 3	(b) $\sqrt{3}$	(c) 9	(d) $\pm 3$
(2)	$\sqrt{8} - \sqrt{2} = \dots\dots\dots$	(a) $\sqrt{6}$	(b) $\sqrt{2}$	(c) 2	(d) 1
(3)	$(\sqrt{8} + \sqrt{2})^2 = \dots\dots\dots$	(a) $\sqrt{10}$	(b) 10	(c) 18	(d) $\sqrt{18}$
(4)	$(\sqrt{7} - \sqrt{5})(\sqrt{7} + \sqrt{5}) = \dots\dots\dots$	(a) 2	(b) 12	(c) $2\sqrt{7}$	(d) $-2\sqrt{5}$
(5)	$\sqrt{\frac{1}{2}} + \sqrt{\frac{1}{2}} = \dots\dots\dots$	(a) 1	(b) $\sqrt{\frac{1}{4}}$	(c) $\sqrt{2}$	(d) $\frac{\sqrt{2}}{2}$
(6)	$\frac{\sqrt{27}}{\sqrt{3}} \div \frac{\sqrt{72}}{\sqrt{2}} = \dots\dots\dots$	(a) $\frac{1}{2}$	(b) 2	(c) -2	(d) 4
(7)	The multiplicative inverse of the number $\sqrt{50}$ is $\dots\dots\dots$	(a) $\frac{\sqrt{2}}{10}$	(b) $\frac{-\sqrt{2}}{10}$	(c) $-5\sqrt{2}$	(d) $5\sqrt{2}$
(8)	If : $x = \frac{\sqrt{6}}{\sqrt{2}}$ , then $x^{-1} = \dots\dots\dots$	(a) $\sqrt{3}$	(b) $\frac{\sqrt{3}}{2}$	(c) $\frac{\sqrt{3}}{3}$	(d) $2\sqrt{3}$

(9) If :  $x = \sqrt{7} + \sqrt{3}$  and  $y = \sqrt{28} + \sqrt{12}$  , then  $x = \dots\dots\dots$

(a)  $y$

(b)  $\frac{1}{2} y$

(c)  $2y$

(d)  $y^2$

[5] If  $x = \frac{\sqrt{50} - \sqrt{18}}{2}$  and  $y = 2 - \sqrt{2}$ , find in the simplest form:

(1)  $x + y = \dots\dots\dots$

(2)  $xy = \dots\dots\dots$

## Sheet (8)

# The two conjugate numbers

If a and b are two positive rational numbers , then each of the two numbers  $(\sqrt{a} + \sqrt{b})$  and  $(\sqrt{a} - \sqrt{b})$  is conjugate to the other one and we find that

- Their sum =  $(\sqrt{a} + \sqrt{b}) + (\sqrt{a} - \sqrt{b}) = 2\sqrt{a}$  = twice the first term.
- Their product =  $(\sqrt{a} + \sqrt{b})(\sqrt{a} - \sqrt{b}) = (\sqrt{a})^2 - (\sqrt{b})^2 = a - b$

The difference:

$$\text{Greater} - \text{smaller} = (\sqrt{a} + \sqrt{b}) - (\sqrt{a} - \sqrt{b}) = 2\sqrt{b} \quad (\text{twice the second term})$$

$$\text{Smaller} - \text{greater} = (\sqrt{a} - \sqrt{b}) - (\sqrt{a} + \sqrt{b}) = -2\sqrt{b} \quad (\text{negative twice the second term})$$

**Example:**  $(\sqrt{2} + \sqrt{5})$  its conjugate is  $(\sqrt{2} - \sqrt{5})$

Their sum 2 of 1 <sup>st</sup> term	G - S 2 of 2 <sup>nd</sup> term	S - G -2 of 2 <sup>nd</sup> term	Their product $(1^{st})^2 - (2^{nd})^2$
$2\sqrt{2}$	$2\sqrt{5}$	$-2\sqrt{5}$	$2 - 5 = -3$

**Exercise (1):**  $(3 - \sqrt{7})$  its conjugate is .....

Their sum	G - S	S - G	Their product

**Exercise (2):**  $(3\sqrt{5} - \sqrt{6})$  its conjugate is .....

Their sum	G - S	S - G	Their product

## Remarks:

- $x^2 - y^2 = (x + y)(x - y)$
- $x^2 + 2xy + y^2 = (x + y)^2$
- $x^2 - 2xy + y^2 = (x - y)^2$

## [1] Choose the correct answer:

(1)	The conjugate of $(\sqrt{3} - \sqrt{5})$ is ..... (a) $\sqrt{5} - 3$ (b) $\sqrt{3} + \sqrt{5}$ (c) $-\sqrt{3} - \sqrt{5}$ (d) $\sqrt{5} - \sqrt{3}$
(2)	$(\sqrt{5} + \sqrt{3})^2 (\sqrt{5} - \sqrt{3})^2 = \dots\dots\dots$ (a) 4 (b) 2 (c) 8 (d) 3
(3)	The number $\frac{4}{3+\sqrt{5}}$ in the simplest form is ..... (a) $3 + \sqrt{5}$ (b) $3 - \sqrt{5}$ (c) $\sqrt{3} + \sqrt{5}$ (d) $3\sqrt{5}$
(4)	$[2, 5] - \{2\} = \dots\dots\dots$ (a) $]2, 5[$ (b) $]2, 5]$ (c) $[2, 5[$ (d) $[2, 5]$
(5)	The conjugate of the number $\frac{1}{\sqrt{3}+\sqrt{2}}$ is ..... (a) $\sqrt{3} - \sqrt{2}$ (b) $\sqrt{3} + \sqrt{2}$ (c) $\frac{1}{\sqrt{3}-\sqrt{2}}$ (d) $-\sqrt{3} - \sqrt{2}$

[2] If  $x = \sqrt{5} + \sqrt{3}$  and  $y = \frac{2}{\sqrt{5}+\sqrt{3}}$ , prove that x and y are conjugate numbers then find the value of  $x^2 + 2xy + y^2$ .

.....

.....

.....

.....

[3] If  $x = \sqrt{5} - \sqrt{2}$  and  $y = \frac{3}{\sqrt{5}-\sqrt{2}}$ , prove that  $x$  and  $y$  are conjugate numbers then find the value of  $x^2 - 2xy + y^2$ .

---

---

---

---

[4] If  $a = \sqrt{3} + \sqrt{2}$  and  $b = \frac{1}{\sqrt{3}+\sqrt{2}}$ , find the value of  $a^2 - b^2$ .

---

---

---

---

[5] If  $x = \frac{4}{\sqrt{7}-\sqrt{3}}$  and  $y = \frac{4}{\sqrt{7}+\sqrt{3}}$ , find the value of  $x^2 y^2$ .

---

---

---

---

[6] If  $x = \sqrt{5} + \sqrt{2}$  and  $y = \sqrt{5} - \sqrt{2}$ , find the value of  $\frac{x+y}{xy-1}$ .

---

---

---

---

[7] If  $x = \frac{2}{\sqrt{5}-\sqrt{3}}$  and  $y = \frac{2}{\sqrt{5}+\sqrt{3}}$ , find the value of  $x^2 - xy + y^2$ .

---

---

---

---

[8] If  $x = \frac{5\sqrt{2}+3\sqrt{5}}{\sqrt{5}}$  and  $y = \frac{2\sqrt{5}-3\sqrt{2}}{\sqrt{2}}$ , find:

(1)  $x^2 + y^2 =$  .....

---

---

(2)  $xy =$  .....

---

---

(3) Prove that:  $\frac{x^2+y^2}{xy} = 38$

---


---

---

---



**[9] Complete:**

- |      |  |
|------|--|
| (1)  | $(\sqrt{7} + \sqrt{3})(\sqrt{7} - \sqrt{3}) = \dots\dots\dots$   |
| (2)  |  If : $x = 3 + \sqrt{2}$ , then its conjugate is ..... and the product of multiplying $x$ by its conjugate is ..... |
| (3)  | The conjugate number of the number $\frac{1}{\sqrt{3} - \sqrt{2}}$ is .....  |
| (4)  | The conjugate number of the number $1 + \frac{7}{\sqrt{7}}$ in the simplest form is .....  |
| (5)  | The multiplicative inverse for $(\sqrt{3} + \sqrt{2})$ in its simplest form is .....   |
| (6)  | If : $x = 2 + \sqrt{5}$ and $y$ is the conjugate number of $x$ , then $(x - y)^2 = \dots\dots\dots$  |
| (7)  | If : $\frac{x}{5 - \sqrt{5}} = 5 + \sqrt{5}$ , then the value of $x$ in its simplest form is .....   |
| (8)  | If : $\frac{1}{x} = \sqrt{5} - 2$ , then the value of $x$ in its simplest form is .....  |
| (9)  | If : $x = \sqrt{3} + 2$ , $y = \sqrt{3} - 2$ , then $(xy, x + y)$ equals .....   |
| (10) | $(\sqrt{2} + \sqrt{3})^{-9} (\sqrt{2} - \sqrt{3})^{-9} = \dots\dots\dots$  |





## Sheet (9)

### Operations on the cube roots

If  $a$  and  $b$  are two real numbers , then

$$\boxed{1} \quad \sqrt[3]{a} \times \sqrt[3]{b} = \sqrt[3]{ab}$$

For example:

$$\bullet \sqrt[3]{3} \times \sqrt[3]{9} = \sqrt[3]{3 \times 9} = \sqrt[3]{27} = 3$$

$$\bullet \sqrt[3]{2} \times \sqrt[3]{-4} = \sqrt[3]{2 \times -4} = \sqrt[3]{-8} = -2$$

$$\boxed{2} \quad \frac{\sqrt[3]{a}}{\sqrt[3]{b}} = \sqrt[3]{\frac{a}{b}} \text{ (where } b \neq 0 \text{)}$$

For example:

$$\bullet \frac{\sqrt[3]{32}}{\sqrt[3]{4}} = \sqrt[3]{\frac{32}{4}} = \sqrt[3]{8} = 2$$

$$\bullet \frac{\sqrt[3]{54}}{\sqrt[3]{-2}} = \sqrt[3]{\frac{54}{-2}} = \sqrt[3]{-27} = -3$$

#### Remarks

\* If  $a$  and  $b$  are two real numbers , then :

$$\boxed{1} \quad \sqrt[3]{a^3 + b^3} \neq a + b, \quad \sqrt[3]{a^3 - b^3} \neq a - b \quad \boxed{2} \quad \sqrt[3]{-a} = -\sqrt[3]{a}$$

$$\boxed{3} \quad a\sqrt[3]{b} = \sqrt[3]{a^3b}$$

For example :  $\bullet 3\sqrt[3]{\frac{1}{9}} = \sqrt[3]{27 \times \frac{1}{9}} = \sqrt[3]{3}$

$$\bullet 8\sqrt[3]{\frac{1}{4}} = 4 \times 2\sqrt[3]{\frac{1}{4}} = 4\sqrt[3]{8 \times \frac{1}{4}} = 4\sqrt[3]{2}$$

$$\boxed{4} \quad \sqrt[3]{\frac{a}{b}} = \sqrt[3]{\frac{a}{b} \times \frac{b^2}{b^2}} = \sqrt[3]{\frac{ab^2}{b^3}} = \frac{1}{b}\sqrt[3]{ab^2}$$

For example :  $\bullet \sqrt[3]{\frac{1}{3}} = \sqrt[3]{\frac{1}{3} \times \frac{9}{9}} = \sqrt[3]{\frac{9}{27}} = \frac{1}{3}\sqrt[3]{9}$

#### Important Remarks

$$\sqrt[3]{16} = \sqrt[3]{8} \times \sqrt[3]{2} = 2\sqrt[3]{2}$$

$$\sqrt[3]{24} = \sqrt[3]{8} \times \sqrt[3]{3} = 2\sqrt[3]{3}$$

$$\sqrt[3]{54} = \sqrt[3]{27} \times \sqrt[3]{2} = 3\sqrt[3]{2}$$

$$\sqrt[3]{81} = \sqrt[3]{27} \times \sqrt[3]{3} = 3\sqrt[3]{3}$$

$$\sqrt[3]{128} = \sqrt[3]{64} \times \sqrt[3]{2} = 4\sqrt[3]{2}$$

$$\sqrt[3]{40} = \sqrt[3]{8} \times \sqrt[3]{5} = 2\sqrt[3]{5}$$

$$\sqrt[3]{250} = \sqrt[3]{125} \times \sqrt[3]{2} = 5\sqrt[3]{2}$$

$$\sqrt[3]{135} = \sqrt[3]{27} \times \sqrt[3]{5} = 3\sqrt[3]{5}$$

[1] Find the result in its simplest form:

$$(1) \quad \sqrt[3]{2} \times \sqrt[3]{32} = \dots\dots\dots$$

$$(2) \quad \frac{\sqrt[3]{72}}{\sqrt[3]{9}} = \dots\dots\dots$$

$$(3) \quad \frac{4\sqrt[3]{-54}}{2\sqrt[3]{-2}} = \dots\dots\dots$$

$$(4) \quad \frac{1}{2}\sqrt[3]{10} \times 6\sqrt[3]{100} = \dots\dots\dots$$

$$(5) \quad \sqrt[3]{\frac{2}{5}} \times \sqrt[3]{\frac{4}{25}} = \dots\dots\dots$$

$$(6) \quad \sqrt[3]{\frac{3}{4}} \div \sqrt[3]{\frac{2}{9}} = \dots\dots\dots$$

[2] Find the result in its simplest form:

$$(1) \quad \sqrt[3]{16} - \sqrt[3]{2} = \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(2) \quad \sqrt[3]{81} + \sqrt[3]{-24} = \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(3) \quad 2\sqrt[3]{54} - 5\sqrt[3]{2} + \sqrt[3]{16} = \dots\dots\dots$$

$$= \dots\dots\dots$$

(4)	$\sqrt[3]{125} - \sqrt[3]{24}$	= .....
		= .....

(5)	$\sqrt[3]{54} + \sqrt[3]{16} - \sqrt[3]{250}$	= .....
		= .....

(6)	$\sqrt[3]{16} - \frac{1}{3}\sqrt[3]{54} + \sqrt[3]{-2}$	= .....
		= .....

(7)	$\sqrt[3]{128} + \sqrt[3]{16} - 2\sqrt[3]{54}$	= .....
		= .....

(8)	$\sqrt[3]{54} \times \sqrt[3]{16} \div (\sqrt[3]{4} \times 6)$	= .....
		= .....



[3] Simplify each of the following:

(1)	$\frac{7}{3}\sqrt{18} + \sqrt[3]{54} - 7\sqrt{2} + \sqrt[3]{16}$	= .....
		= .....

(2)	$\sqrt{27} + \frac{1}{3}\sqrt[3]{27} - 9\sqrt{\frac{1}{3}} - 1$	= .....
		= .....

(3)	$\sqrt[3]{-16} + \frac{14}{\sqrt{7}} - \sqrt{28} + \sqrt[3]{54}$	= .....
		= .....

$$(4) \quad \sqrt[3]{18} + \sqrt[3]{54} - \frac{\sqrt[3]{216}}{\sqrt[3]{12}} - \sqrt[3]{16} = \dots\dots\dots$$

$$= \dots\dots\dots$$

$$(5) \quad 5\sqrt{2} - \frac{1}{2}\sqrt{200} + (\sqrt[3]{5} \times \sqrt[3]{25}) = \dots\dots\dots$$

$$= \dots\dots\dots$$

[4]

 If  $a = \sqrt[3]{5} + 1$  ,  $b = \sqrt[3]{5} - 1$  Find the value of the following :

**1**  $(a - b)^5$

**2**  $(a + b)^3$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

$$= \dots\dots\dots$$

[5]

If  $x = 3 + \sqrt[3]{6}$  ,  $y = 3 - \sqrt[3]{6}$  Find the value of :  $\left(\frac{x-y}{x+y}\right)^3$

$$\dots\dots\dots$$


$$\dots\dots\dots$$

$$\dots\dots\dots$$

$$\dots\dots\dots$$

$$\dots\dots\dots$$

**[6] Complete:**

(1)	$\sqrt[3]{\frac{2}{3}} \times \sqrt[3]{-12} = \dots\dots\dots$
(2)	$\sqrt[3]{3} \times \sqrt[3]{9} = \sqrt{\dots\dots\dots}$
(3)	$\sqrt[3]{54} - \sqrt[3]{-16} = \sqrt[3]{\dots\dots\dots}$
(4)	The conjugate of the number $\frac{2\sqrt{5} - 3\sqrt{2}}{\sqrt{2}}$ is $\dots\dots\dots$
(5)	$[0, 5] - [0, 3] = \dots\dots\dots$ 
(6)	$\sqrt[3]{54} - \sqrt[3]{2} = \dots\dots\dots$
(7)	If : $x = \sqrt[3]{2} + 1$ and $y = \sqrt[3]{2} - 1$ , then $(x + y)^3 = \dots\dots\dots$
(8)	$2\sqrt{\frac{1}{2}} - \sqrt{2} = \dots\dots\dots$
(9)	The number $-\sqrt{11}$ is included between the two consecutive integers $\dots\dots\dots$ and $\dots\dots$

**[7] Choose the correct answer:**

(1)	$\sqrt[3]{54} + \sqrt[3]{-2} = \dots\dots\dots$ (a) $\sqrt[3]{52}$ (b) $\sqrt[3]{2}$ (c) $2\sqrt[3]{2}$ (d) $4\sqrt[3]{2}$
(2)	$\sqrt[3]{-64} + \sqrt{16} = \dots\dots\dots$ (a) zero                      (b) 8                      (c) - 8                      (d) $\pm 8$
(3)	$\frac{\sqrt[3]{16}}{\sqrt[3]{2}} = \dots\dots\dots$ (a) 8                      (b) -2                      (c) 2                      (d) $2\sqrt[3]{2}$





(4)	$\sqrt[3]{2} + \sqrt[3]{2} = \dots\dots\dots$ (a) $\sqrt[3]{2}$ (b) $\sqrt[3]{4}$ (c) $\sqrt[3]{8}$ (d) $\sqrt[3]{16}$
(5)	$\sqrt[3]{\frac{2}{9}} = \dots\dots\dots$ (a) $\frac{\sqrt[3]{6}}{3}$ (b) $\sqrt[3]{\frac{1}{6}}$ (c) $\sqrt[3]{6}$ (d) $\sqrt[3]{2}$
(6)	If : $X = ]-\infty, 0[$ , then $\tilde{X} = \dots\dots\dots$ (a) $\mathbb{R}_+$ (b) $[0, \infty[$ (c) $]-\infty, 0]$ (d) $\mathbb{R}_-$
(7)	The multiplicative inverse of the number $\sqrt{\frac{3}{2}}$ is $\dots\dots\dots$ (a) $\frac{2}{3}\sqrt{2}$ (b) $\frac{3\sqrt{2}}{2}$ (c) $\frac{\sqrt{6}}{3}$ (d) $-\sqrt{\frac{3}{2}}$
(8)	The irrational number in the following numbers is $\dots\dots\dots$ (a) $\sqrt{\frac{1}{4}}$ (b) $\sqrt[3]{8}$ (c) $\sqrt{\frac{4}{9}}$ (d) $2\sqrt{2}$
(9)	$]-1, 3] \cap [-3, -1] = \dots\dots\dots$ (a) $\emptyset$ (b) $\{-3\}$ (c) $\{-1\}$ (d) $\{3\}$
(10)	If : $x = \sqrt{3} + \sqrt{2}$ and $xy = 1$ , then $y = \dots\dots\dots$ (a) $\sqrt{2} - \sqrt{3}$ (b) $\sqrt{3} + \sqrt{2}$ (c) $\sqrt{3} - \sqrt{2}$ (d) 1



## Sheet (10)

### Applications on the real numbers

In the following , we will summarize the previous rules of areas and volumes of some solids :

The solid	The lateral area	The total area	The volume
The cube 	$4l^2$	$6l^2$	$l^3$
The cuboid 	$2(x+y) \times z$	$2(xy + yz + zx)$	$xyz$
The cylinder 	$2\pi r h$	$2\pi r h + 2\pi r^2$ $= 2\pi r (h + r)$	$\pi r^2 h$
The sphere 	—	$4\pi r^2$	$\frac{4}{3}\pi r^3$

### THE CUBE

[1] A cube with volume  $125 \text{ cm}^3$ . Find its total area and its lateral area.



[2] A cube whose lateral area is  $36 \text{ cm}^2$ . Find its total area, and its volume.

---

---

---

[3] The perimeter of one face of a cube is  $12 \text{ cm}$ . Find its volume, and its lateral area.

---

---

---

[4] The sum of lengths of all edges of a cube is  $60 \text{ cm}$ . Find its volume and its total area.

---



---

---

[5] Complete:

- |     |   |
|-----|---|
| (1) | If the edge length of a cube is $5 \text{ cm}$ , then its volume = ..... $\text{cm}^3$  |
| (2) | The edge length of a cube is $4 \text{ cm}$ , then its total area = ..... $\text{cm}^2$ |
| (3) | The lateral area of a cube whose edge length is $l \text{ cm}$ . = ..... $\text{cm}^2$  |
| (4) | The cube whose volume $l^3 \text{ cm}^3$ , its total area = ..... $\text{cm}^2$         |
| (5) | The cube whose edge length is $2l$ , then its volume = ..... $\text{cm}^3$              |

**[6] Choose the correct answer:**

(1)	The volume of a cube is $1 \text{ cm}^3$ , then the sum of its edge lengths = ..... cm. (a) 1                      (b) 6                      (c) 8                      (d) 12
(2)	 The volume of a cube is $64 \text{ cm}^3$ , then its lateral area = ..... $\text{cm}^2$ . (a) 4                      (b) 8                      (c) 64                      (d) 96
(3)	If the total area of a cube is $96 \text{ cm}^2$ , then the area of one face = ..... $\text{cm}^2$ . (a) 16                      (b) 64                      (c) 24                      (d) 48
(4)	If the area of the six faces of a cube = $54 \text{ cm}^2$ , then its volume = ..... $\text{cm}^3$ . (a) 54                      (b) 44                      (c) 72                      (d) 27
(5)	If the volume of a cube = $64 \text{ cm}^3$ , then the length of a diagonal of one face = ..... cm. (a) 16                      (b) $4\sqrt{2}$ (c) 32                      (d) 64
(6)	The volume of a cube is $5 \text{ cm}^3$ . If the edge length became twice the first, then its volume = ..... $\text{cm}^3$ . (a) 10                      (b) 20                      (c) 30                      (d) 40
(7)	 The edge length of a cube whose volume is $2\sqrt{2} \text{ cm}^3$ = ..... cm. (a) $\sqrt{2}$ (b) 2                      (c) 8                      (d) 1.5

## THE CUBOID

**[1] The dimensions of the base of a cuboid are 9 cm and 10 cm and its height is 5 cm. Find its volume, its lateral area and its total area.**

.....

.....

.....


[2] The height of a cuboid is 4 cm and its base is a square of side length 5 cm. Find its volume, its lateral area and its total area.

[3] The lateral area of a cuboid is 480 cm<sup>2</sup> and its base is in the shape of a square whose side length is 10 cm. Calculate its height.

## THE CIRCLE

If  $M$  is a circle with radius length  $r$ , then :

- 1 The circumference of the circle =  $2 \pi r$  length unit.
- 2 The area of the circle =  $\pi r^2$  square unit.



[1] A circle is of radius length 10.5 cm. Find each of its circumference and its area.  $\left(\pi = \frac{22}{7}\right)$

[2] The area of a circle is  $25\pi \text{ cm}^2$ . Calculate its circumference in terms of  $\pi$ .

---

---

---


[3] The area of a circle is  $154 \text{ cm}^2$ . Find its circumference and its diameter length.  $\left(\pi = \frac{22}{7}\right)$

---

---

---

## THE RIGHT CIRCULAR CYLINDER



- 1 The lateral area of the cylinder =  $2 \pi r h$  square unit.
- 2 The total area of the cylinder = the lateral area of the cylinder + twice the area of the base  
 $= 2 \pi r h + 2 \pi r^2$  square unit.
- 3 The volume of the cylinder = the area of the base  $\times$  height =  $\pi r^2 h$  cube unit.

Consider  $\pi = \frac{22}{7}$  if there are not any other values given.

- (1) A right circular cylinder, the radius length of its base is 14 cm. and its height is 20 cm.  
 Find the volume and the total area of the cylinder. « 12320 cm<sup>3</sup>, 2992 cm<sup>2</sup>. »

---

---

(2)

Find the lateral area for a right circular cylinder of volume  $924 \text{ cm}^3$ , and of a height 6 cm.

«  $264 \text{ cm}^2$ . »


(3)

 Find the total area of a right circular cylinder of volume  $7536 \text{ cm}^3$  and its height is 24 cm.

( $\pi = 3.14$ )

«  $2135.2 \text{ cm}^2$ . »

(4)

 Find the height of a right circular cylinder whose height is equal to its base radius length and its volume is  $72 \pi \text{ cm}^3$ .

«  $2\sqrt[3]{9} \text{ cm}$ . »

## THE SPHERE

**1** The area of the sphere =  $4 \pi r^2$  square unit.

**2** The volume of the sphere =  $\frac{4}{3} \pi r^3$  cube unit.

Consider  $\pi = \frac{22}{7}$  if there are not any other values given.

(1)

Find the volume and the surface area of a sphere if the length of its diameter is 4.2 cm.

«  $38.808 \text{ cm}^3$ ,  $55.44 \text{ cm}^2$ . »

(2) The volume of a sphere is  $4188 \text{ cm}^3$ . Find its radius length. ( $\pi = 3.141$ ) « 10 cm. »


.....

.....

(3) The volume of a sphere =  $\frac{500}{3} \pi \text{ cm}^3$ . Find the length of its diameter.

.....

.....

(4)  The volume of a sphere is  $562.5 \pi \text{ cm}^3$ . Find its surface area in terms of  $\pi$  «  $225 \pi$  »

.....

.....

## General Revision on Applications on the Real Numbers

### [1] Complete:

(1) The sphere whose volume =  $36 \pi \text{ cm}^3$  has a radius length = ..... cm.


(2) A right circular cylinder, its volume is  $343\pi \text{ cm}^3$ . If its height equals its base radius length, then its height = ..... cm.

(3) The volume of a cube is  $64 \text{ cm}^3$ , then its total area is .....  $\text{cm}^2$ .

### [2] Choose the correct answer:

(1) The circle whose radius length =  $\sqrt{14} \text{ cm}$  has an area = .....  $\text{cm}^2$   
 (a)  $14 \pi$  (b)  $2\sqrt{14} \pi$  (c) 14 (d)  $2\sqrt{14}$

(2)  $]2, 4[ \cup \{2, 4\} = \dots\dots\dots$   
 (a)  $\emptyset$  (b)  $\{2, 4\}$  (c)  $[2, 4]$  (d)  $]2, 4[$

(3)	The right circular cylinder whose base radius length = 3 cm. and its height = 5 cm. its volume = ..... $\text{cm}^3$ (a) $15 \pi$ (b) $75 \pi$ (c) $45 \pi$ (d) $\frac{3}{5} \pi$
(4)	If : $-\sqrt[3]{25} = \sqrt[3]{x}$ , then $x = \dots\dots\dots$ (a) $-5$ (b) $-25$ (c) $-125$ (d) $125$
(5)	 The volume of the sphere whose diameter length is 6 cm. = ..... $\text{cm}^3$ (a) 288 (b) $12 \pi$ (c) $36 \pi$ (d) $288 \pi$
(6)	If the volume of a sphere = $\frac{9}{16} \pi \text{ cm}^3$ , then its radius length = ..... cm. (a) 3 (b) $\frac{4}{3}$ (c) $\frac{3}{4}$ (d) $\frac{1}{3}$
(7)	If the surface area of a sphere is $9 \pi \text{ cm}^2$ , then its diameter length = ..... cm. (a) 9 (b) 3 (c) 1.5 (d) 6
(8)	If three quarters of volume of a sphere equals $8 \pi \text{ cm}^3$ , then the length of its radius equals ..... cm. (a) 64 (b) 8 (c) 4 (d) 2



**Sheet (11)**  
**Solving equations and and inequalities**  
**of the first degree in one variable in  $\mathbb{R}$**

[1] Find the solution set for each of the following equations in  $\mathbb{R}$ , then graph the solution on the number line:

(1)  $x + 5 = 0$

.....  
 .....



(2)  $5x + 6 = 1$

.....  
 .....



(3)  $2x + 4 = 3$

.....  
 .....



(4)  $2x - 3 = 4$

.....  
 .....



(5)  $\sqrt{5}x - 1 = 4$

.....  
 .....



(6)  $x - 1 = \sqrt{3}$

.....  
 .....



(7)  $\sqrt{3}x - 1 = 2$

.....  
 .....



(8)  $7x - \sqrt{7} = 6\sqrt{7}$

.....  
 .....



(9)  $x - \sqrt{5} = 1$

.....  
.....



(10)  $x - \sqrt{3} = 1$

.....  
.....



[2] Find the solution set for each of the following inequalities in  $\mathbb{R}$  in the form of interval, then graph the solution on the number line:

(1)  $2x > 6$

.....  
.....



(2)  $-7x \geq -14$

.....  
.....



(3)  $x + 3 \leq 5$

.....  
.....



(4)  $5 - x > 3$

.....  
.....



(5)  $2x + 5 \geq 3$

.....  
.....



(6)  $1 - 5x < 6$

.....  
.....



(7)  $\frac{1}{2}x + 1 \leq 2$

.....  
.....



(8)  $3 - 2x \leq 7$

.....  
.....



(9)  $3 < x + 2 \leq 6$

.....  
.....



(10)  $-5 < x + 3 < 9$

.....  
.....



(11)  $-3 \leq -x \leq 3$

.....  
.....



(12)  $1 < 5 - x \leq 3$

.....  
.....



(13)  $\sqrt[3]{-8} \leq x + 1 \leq \sqrt{9}$

.....  
.....



(14)  $5 < 3 - x \leq 3^2$

.....  
.....



(15)  $|-3| < 2x - 1 < 5$

.....  
.....



(16)  $3x < 2x + 4$

.....  
.....



(17)  $7x - 9 \geq 4x$

.....  
.....



(18)  $5x - 3 < 2x + 9$

.....  
.....



(19)  $x + 3 \geq 2x \geq x - 2$

.....  
.....



(20)  $4x \leq 5x + 2 \leq 4x + 3$

.....  
.....



(21)  $x - 1 < 3x - 1 \leq x + 1$

.....  
 .....



(22)  $2x - 1 < 3x - 2 \leq 2x + 1$

.....  
 .....



(23)  $1 \leq 3 - 2x < 5$

.....  
 .....



(24)  $2x - 1 \geq 7$

.....  
 .....



(25)  $-7 \leq 3x + 2 < 11$

.....  
 .....



(26)  $x + 1 \leq 2x - 3 < x + 4$

.....  
 .....



(27)  $-2 < 3x + 7 \leq 10$

.....  
 .....



(28)  $-3 < 2x - 3 \leq \sqrt[3]{125}$

.....  
 .....



(29)  $3 \leq 2x - 1 < 11$

.....  
 .....



(30)  $6 < 2x + 4 \leq 10$

.....  
 .....



(31)  $5 < 2x - 3 \leq 11$

.....  
 .....



(32)  $2x + 4 \geq 3x - 3 > 2x + 1$

.....  
 .....



**[1] Complete:**

(1)	If $x - 3 \geq 0$ , then $x$ .....
(2)	If $5x < 15$ , then $x$ .....
(3)	If $1 - x > 4$ , then $x$ .....
(4)	If $-2x \leq 3$ , then $x$ .....
(5)	If $\sqrt{2}x \leq 4$ , then $x$ .....
(6)	The S.S. of the inequality $-5 \leq -x < 2$ in $\mathbb{R}$ is .....
(7)	If $-3 < x < 3$ where $x \in \mathbb{R}$ , then $2x \in ] - 6 , \dots [$

**[2] Choose the correct answer:**

(1)	The S.S. of the inequality : $x + 3 < 3$ in $\mathbb{R}$ is ..... (a) $] - \infty , 0[$ (b) $] - \infty , 0]$ (c) $[ 0 , \infty [$ (d) $] 0 , \infty [$
(2)	The S.S. of the inequality : $1 > x - 5 > -1$ in $\mathbb{R}$ is ..... (a) $[ 4 , 6]$ (b) $] 4 , 6[$ (c) $] 4 , 6]$ (d) $[ 4 , 6[$
(3)	If $x > 5$ , then $-x$ ..... (a) $< -9$ (b) $\geq -5$ (c) $< -5$ (d) $> -5$
(4)	If $-2 < x < 2$ , then $2x + 3$ belongs to ..... (a) $[-1 , 7[$ (b) $] -1 , 5[$ (c) $] -1 , 7[$ (d) $] -4 , 6[$

## Sheet (12)

# Relation between two variables

- (1) Complete the following ordered pairs which satisfy the relation :  $y = 3x - 1$   
 (5 , .....), (2 , .....), (0 , .....), (- 3 , .....)
- (2) Show which of the following ordered pairs satisfy the relation :  $y - 4x = 7$   
 1 (1 , 2)                      2 (3 , - 5)                      3 (- 1, 3)
- (3) Find four ordered pairs satisfying the relation  $y = 2x - 1$  and represent it. ....



- (4) Find three ordered pairs that satisfy the relation  $x + 2y = 6$ , then represent it graphically.....



- (5) Graph the relation  $y = 2x - 3$ .....





(6) Represent graphically the relation  $x + y = 2$





(7) Represent the relation  $2y - x = 2$  graphically



**[8] Complete:**

(1)	If (2,3) satisfies the relation $x + y = k$ , then $k = \dots\dots\dots$
(2)	If (k,2k) satisfies the relation $x + y - 15 = 0$ , then $k = \dots\dots\dots$
(3)	If (-2,k) satisfies the relation $2x + 3y = 35$ , then $k = \dots\dots\dots$

**[9] Choose the correct answer:**

(1)	If (2 , - 5) satisfies the relation : $3x - y + c = 0$ , then $c = \dots\dots\dots$ (a) 1                                      (b) - 1                                      (c) 11                                      (d) - 11
(2)	 Which of the following ordered pairs satisfies the relation : $2x + y = 5$ ? (a) (- 1 , 3)                              (b) (1 , 3)                                      (c) (3 , 1)                                      (d) (2 , 2)
(3)	 (3 , 2) does not satisfy the relation ..... (a) $y + x = 5$ (b) $3y - x = 3$ (c) $y + x = 7$ (d) $x - y = 1$
(4)	The relation : $3x + 8y = 24$ is represented by a straight line intersecting y-axis at the point ..... (a) (0 , 8)                                      (b) (8 , 0)                                      (c) (0 , 3)                                      (d) (3 , 0)
(5)	The relation $2x + 7y = 14$ is represented by a straight line intersecting x-axis at the point ..... (a) (2 , 0)                                      (b) (0 , 2)                                      (c) (7 , 0)                                      (d) (0 , 7)
(6)	If : (2 k , 3 k) satisfies the relation $x + y = 15$ , then $k = \dots\dots\dots$ (a) 5    (b) 3    (c) - 5    (d) - 3

# Sheet (13)

## Slope of straight line

If a point moves on a straight line  $L$  from the location  $A (x_1, y_1)$  to the location

$B (x_2, y_2)$ , then :


The change in the  $x$ -coordinates =  $x_2 - x_1$

It is called (the horizontal change).

The change in the  $y$ -coordinates =  $y_2 - y_1$

It is called (the vertical change).

The ratio of the change in the  $y$ -coordinates to the change in the  $x$ -coordinates is called the slope of the straight line ( $m$ ).



The slope of the straight line =  $\frac{\text{the change in } y\text{-coordinates}}{\text{the change in } x\text{-coordinates}} = \frac{\text{the vertical change}}{\text{the horizontal change}}$

i.e.  $S = \frac{y_2 - y_1}{x_2 - x_1}$ , where  $x_1 \neq x_2$ ,  $S$  is undefined if  $x_1 = x_2$

**1** Classify the slope of the straight line in each of the following figures showing whether it is (positive – negative – zero – undefined)




Fig. (1)




Fig. (2)




Fig. (3)




Fig. (4)

**[2] Find the slope of the straight line passing through the two points in each of the following:**

(1) A (1 , 3) , B (3 , 4)

(2) A (1 , 2) , B (5 , 0)

(3) A (2 , - 1) , B (4 , - 1)

(4) A (5 , 2) , B (5 , 4)

(5) A (3,5) , B (5,-1)

(6) A (-1,3) , B (2,4)

(7) A (1,3) , B (2,1)

(8) A (1,3) , B (2,3)

**[3]**

 **In the opposite figure :**


ABC is a triangle. Complete by using one of the following words  
(positive , negative , zero , undefined)

1 The slope of  $\overleftrightarrow{AB}$  is .....

2 The slope of  $\overleftrightarrow{BC}$  is .....

3 The slope of  $\overleftrightarrow{AO}$  is .....

4 The slope of  $\overleftrightarrow{AC}$  is .....




**[4] Complete:**


(1)	The slope of any horizontal straight line equals .....
(2)	The slope of any straight line parallel to y-axis is .....
(3)	The straight line whose slope = zero is parallel to .....
(4)	If A , B and C are collinear then the slope of $\overleftrightarrow{AB}$ = the slope of .....
(5)	The slope of the straight line $\overleftrightarrow{AB}$ where A (2 , 3) and B (0 , 4) is .....
(6)	If the slope of the straight line which passes through the two points (1 , 3) , (3 , k) equals 3 , find the value of k « 9 »
(7)	If the slope of the straight line which passes through the two points (3 , c) and (5 , - 2) equals - 3 , find the value of c « 4 »

**[4] Prove that the points A(2,5) , B(0,1) and C(5,12) are collinear.**

# Real life applications on the slope

- 1**  An irrigation machine consumes 2.47 litres of diesel to work for 3 hours. If the machine works for 10 hours , how many litres of diesel will the machine consume ? «  $8\frac{7}{30}$  litres »


- 2** A car moves with uniform velocity such that it covers 180 km. per 3 hours. If the car moves for 5 hours , what is the covered distance ? « 300 km. »

- 3**  The opposite figure represents the motion of a bicycle measured from a constant point. Find the regular speed of the bicycle during :


(1) The first three hours.

(2) The next four hours.


Find the total distance covered by the bicycle.



- 4** The opposite graph represents the motion of a car moving with uniform velocity. Determine the velocity of the car.







- 5  The opposite figure shows the capital change of a company during 8 years :


- (1) Find the slope of each of  $\overrightarrow{AB}$  ,  $\overrightarrow{BC}$  and  $\overrightarrow{CD}$   
What is the meaning of each ?  
(2) Find the starting capital of the company.

« 10 , 0 , - 5 , 20 thousand pounds »



- 6  The opposite figure shows the relation between the height of a person (in cm.) and his age (in years) :


- (1) Find the slope of each of  $\overrightarrow{AB}$  ,  $\overrightarrow{BC}$  and  $\overrightarrow{CD}$   
What is the meaning of each ?  
(2) Calculate the difference between the height of this person when he was 8 years old and his height when he was 30 years old.



«  $9\frac{3}{8}$  , 5 , 0 , 50 cm. »

- 7 The opposite graph shows the relation between the distance in km. and the time (t) in hours for a bicycle which moved between two towns A and B going and returning back. Answer the following :

- (1) What is the uniform velocity during the going trip?  
(2) What is the average velocity during returning back ?  
(3) What is the meaning of the horizontal line segment in the graph ?




« 20 km./hr. , 12 km./hr. »



- 8** A farmer wanted to complete digging a well in his farm. He rented a digging machine. The opposite graph shows the depth of the well (d) in metres after time (t) in hours , find :

- (1) The depth of the well before beginning digging.
- (2) The depth of the well after finishing digging.
- (3) The total time which the machine took in digging the well.
- (4) The average of depth of the well which the machine digs within the first five hours.
- (5) The average of the depth of the well within the last two hours of digging.




« 5 m. , 40 m. , 10 hr. , 4.5 m./hr. , 6.25 m./hr. »

- 9** Magdi filled the tank of his car by fuel. The opposite figure represents the relation between the time (t) in hours and the amount of remained fuel in the tank (y) in litres :

- (1) What is the greatest capacity of the tank ?
- (2) When will the tank become empty ?
- (3) What is the amount of remained fuel after 15 hours ?
- (4) What is the range of consumption of fuel in each hour ?

« 70 L , 30 hr. , 35 L ,  $2\frac{1}{3}$  L /hr. »




10

Bassim drove his car from the city A to the city B  
The opposite graph shows the relation between the distance  $d$  in km. and the time  $t$  in hours.

**Answer the following :**

- (1) What is the uniform velocity of the car of Bassim ?
- (2) Find the distance between the car and the point 0 after three hours from the moment of beginning.




« 75 km./hr. , 275 km. »

11

**A person read a book.**

The opposite graph shows the relation between the time ( $t$ ) in hours and the number of remained pages ( $y$ ) :

- (1) How many pages are remained in the beginning ?
- (2) Find the rate of reading pages per hour.
- (3) When does this person finish reading this book ?



« 100 pages , 20 page/hr. , after 5 hours »



## Sheet (14)

# The ascending and descending cumulative frequency table and their graphical representation

**Example 1**

The following frequency table shows the weekly wages in pounds of 50 workers in one factory :

Sets of wages	54 –	58 –	62 –	66 –	70 –	Total
No. of workers (Frequency)	5	12	22	7	4	50

Form the ascending cumulative frequency table and represent it graphically , then find :

- 1 The number of workers whose weekly wages are less than 60 pounds.
- 2 The percentage of the number of workers whose weekly wages are less than 60 pounds.

**Solution**

- Form the ascending cumulative frequency table as follows :

The upper boundaries of sets	Frequency
Less than 54	
Less than 58	
Less than 62	
Less than 66	
Less than 70	
Less than 74	

Sets of wages	54 –	58 –	62 –	66 –	70 –
Number of workers (Frequency)	5	12	22	7	4

Less than 54 =   
 Less than 58 = 5 + 0 =   
 Less than 62 = 5 + 12 =   
 Less than 66 = 5 + 12 + 22 =   
 Less than 70 = 5 + 12 + 22 + 7 =   
 Less than 74 = 5 + 12 + 22 + 7 + 4 =


The ascending cumulative frequency table.

**Notice that :** The ascending cumulative frequency begins with zero and ends at the total frequency. To represent the ascending cumulative frequency table graphically , do as follows :

- 1 Specialize the horizontal axis for sets and the vertical axis for the ascending cumulative frequency.
- 2 Choose a suitable scale to represent data on the vertical axis so that it contains the ascending cumulative frequency easily.
- 3 Represent the ascending cumulative frequency of each set , then draw the graph (the curve) such that it passes through the points which we located as shown in the opposite figure.

• From the graph , we find that :

- 1 The number of workers whose weekly wages are less than 60 pounds = 10 workers.
- 2 The percentage of the number of workers whose weekly wages are less than 60 pounds =  $\frac{10}{50} \times 100\% = 20\%$





### Example 2

The following frequency table shows the weekly wages of 50 workers in one factory :

Sets of wages	54 –	58 –	62 –	66 –	70 –	Total
No. of workers (Frequency)	5	12	22	7	4	50

Form the descending cumulative frequency table and represent it graphically , then find :

- 1 The number of workers whose weekly wages are 60 pounds or more.
- 2 The percentage of the number of workers whose weekly wages are 60 pounds or more.

### Solution

- Form the descending cumulative frequency table as follows :

Sets of wages	54 –	58 –	62 –	66 –	70 –
Number of workers (Frequency)	5	12	22	7	4

$$\begin{aligned}
 54 \text{ and more} &= 5 + 12 + 22 + 7 + 4 = \\
 58 \text{ and more} &= 12 + 22 + 7 + 4 = \\
 62 \text{ and more} &= 22 + 7 + 4 = \\
 66 \text{ and more} &= 7 + 4 = \\
 70 \text{ and more} &= \\
 74 \text{ and more} &=
 \end{aligned}$$

The lower boundaries of sets	Frequency
54 and more	
58 and more	
62 and more	
66 and more	
70 and more	
74 and more	

The descending cumulative frequency table


**Notice that :** The descending cumulative frequency begins with the total frequency and ends with zero.

- To represent this table graphically , follow the same previous steps in the ascending cumulative frequency table to get the opposite graph.

- From the graph , we find that :

- 1 The number of workers whose weekly wages are 60 pounds or more = 40 workers.

- 2 The percentage of those workers =  $\frac{40}{50} \times 100\% = 80\%$



The descending cumulative frequency curve

# Sheet (15)

## The Mean

### Remember that :

To calculate the mean of a set of values, do as follows :

- 1** Find the sum of these values.      **2** Divide this sum by the number of these values

i.e. The mean of a set of values =  $\frac{\text{The sum of values}}{\text{Number of values}}$

### For example:

If the marks of 5 students are 25 , 23 , 21 , 22 , 24

, then the mean of marks =  $\frac{25 + 23 + 21 + 22 + 24}{5} = 23$  marks.

Notice that :  $23 \times 5 = 115$

, the sum of marks of the 5 students =  $25 + 23 + 21 + 22 + 24 = 115$

i.e. The mean is the value which is given to each item of a set , then the sum of these new values is the same sum of the original values.

### Finding the mean of data from the frequency table with sets

**Example** The following table shows the distribution of the marks of 50 students in mathematics :

Sets	10 –	20 –	30 –	40 –	50 –	Total
Frequency	8	12	14	9	7	50

Find the mean of these marks.

- 1** Determine the centres of sets according to the rule :

The centre of a set =  $\frac{\text{the lower limit} + \text{the upper limit}}{2}$

, then the centre of the first set =  $\frac{10 + 20}{2} = 15$

, the centre of the second set =  $\frac{20 + 30}{2} = 25$  ... and so on.

Since the lengths of the subsets are equal and each of them = 10 therefore we consider the upper limit of the last set = 60

, then its centre =  $\frac{50 + 60}{2} = 55$



**2 Form the vertical table :**

Set	Centre of the set « $X$ »	Frequency « $f$ »	$X \times f$
10 –		8	
20 –		12	
30 –		14	
40 –		9	
50 –		7	
<b>Total</b>		50	

**3** The mean =  $\frac{\text{The sum of } (X \times f)}{\text{The sum of } f} =$

## Sheet (16)

# The Median

**Remember that**

The median is the middle value in a set of values after arranging it ascendingly or descendingly, such that the number of values which are less than it is equal to the number of values which are greater than it.

- To find the median of a set of values , we do as follows :

### We arrange the values ascendingly or descendingly

**If the values number is odd , then**

The median is the value lying in the middle exactly.

**For example :**

If the values are  
42 , 23 , 17 , 30 and 20

We arrange them ascendingly as follows

17 , 20 , 23 , 30 , 42

The median = 23

**If the values number is even , then**

The median

=  $\frac{\text{The sum of the two values lying in the middle}}{2}$

**For example :**

If the values are  
27 , 13 , 23 , 24 , 13 , 21

We arrange them ascendingly as follows

13 , 13 , 21 , 23 , 24 , 27

The median =  $\frac{21 + 23}{2} = 22$

## Finding the median of a frequency distribution with sets graphically

**Example** The following table shows the frequency distribution of marks of 50 students in math exam :

Sets of marks	0 –	10 –	20 –	30 –	40 –	50 –	Total
Number of students	2	5	8	19	14	2	50

Find the median mark of the student.



### Solution

Using the ascending cumulative frequency curve :

The upper boundaries of sets	Frequency
Less than 0	
Less than 10	
Less than 20	
Less than 30	
Less than 40	
Less than 50	
Less than 60	

Using the descending cumulative frequency curve :

The lower boundaries of sets	Frequency
0 and more	
10 and more	
20 and more	
30 and more	
40 and more	
50 and more	
60 and more	



$\therefore$  The order of the median =  $\frac{50}{2} = 25$

$\therefore$  From the two previous graphs , the median = 36 approximately



# Sheet (17)

## The Mode

### Remember that



The mode of a set of values is the most common value in the set, or in other words, it is the value which is repeated more than any other values.

### For example :

The mode of the set of the values : 7 , 3 , 4 , 1 , 7 , 9 , 7 , 4 is 7

### Finding the mode for a frequency distribution with equal sets in range.

The following is an example which shows how to find the mode of a frequency distribution with sets.

### Example

The following is the frequency distribution of marks of 100 pupils in one of the exams :


Set of marks	10 –	20 –	30 –	40 –	50 –	Total
Number of pupils	16	24	30	20	10	100

Find the mode mark for these pupils.

### Solution

We can find the mode of that distribution graphically using the histogram as follows :

- 1 Draw two orthogonal axes : one of them is horizontal and the other is vertical to represent the frequency of each set.
- 2 Divide the horizontal axis into a number of equal parts with a suitable drawing scale to represent the sets.
- 3 Divide the vertical axis into a number of equal parts with a suitable drawing scale to represent the greatest frequency in the sets.
- 4 Draw a rectangle whose base is set (10 –) and its height equals the frequency (16)
- 5 Draw a second rectangle adjacent to the first one whose base is set (20 –) and its height equals the frequency (24)



**[1] Choose the correct answer:**

<b>(1)</b>	The arithmetic mean of : 3 , 10 , 2 is ..... (a) 10 (b) 5 (c) 3 (d) 6
<b>(2)</b>	The mean of the values : 2 , 5 , 4 , 5 is ..... (a) 4 (b) 5 (c) 16 (d) 8
<b>(3)</b>	The mean of the values : 2 , 8 , 6 , 4 is ..... (a) 2 (b) 5 (c) 4 (d) 6
<b>(4)</b>	The arithmetic mean of : 3 , 7 , 28 , 52 , 10 = ..... (a) 17 (b) 19 (c) 20 (d) 27
<b>(5)</b>	The arithmetic mean of the values : 19 , 32 , 21 , 6 , 12 is ..... (a) 90 (b) 32 (c) 18 (d) 6
<b>(6)</b>	The mean of the values : 7 , 15 , 19 , 14 and 15 is ..... (a) 14 (b) 15 (c) 16 (d) 17
<b>(7)</b>	The arithmetic mean of the values : 30 , 23 , 25 , 30 , 22 is ..... (a) 22 (b) 23 (c) 24 (d) 26
<b>(8)</b>	If the arithmetic mean of the values : 27 , 8 , 16 , 24 , 6 and k is 14 , then k = ..... (a) 3 (b) 6 (c) 27 (d) 84
<b>(9)</b>	If the mean of marks of 5 pupils is 20 , then the total of their marks = ..... (a) 4 (b) 15 (c) 25 (d) 100
<b>(10)</b>	If the sum of 5 numbers equals 30 , then the arithmetic mean of these numbers is ..... (a) 150 (b) 6 (c) 18 (d) 72
<b>(11)</b>	The set which its lower boundary is 2 and its upper boundary is 6 , then its centre is ..... (a) 2 (b) 6 (c) 4 (d) 8
<b>(12)</b>	The lowest limit of a set is 4 and the other limit is 8 , then its centre is ..... (a) 2 (b) 4 (c) 6 (d) 8



(13)	If the lower limit of a set is 6 and the upper limit is 10 , then its centre is ..... (a) 4 (b) 6 (c) 10 (d) 8
(14)	If the upper limit of a set is 19 and the lower limit of the same set is 11 , then centre is ..... (a) 10 (b) 15 (c) 20 (d) 30
(15)	If the lowest boundary of a set is 10 and the upper boundary is $X$ and its centre then $X =$ ..... (a) 10 (b) 15 (c) 20 (d) 30
(16)	If the lower limit of a set is 18 and its centre is 20 , then its length is ..... (a) 2 (b) 19 (c) 22 (d) 4
(17)	The arithmetic mean of the values : $3 - a$ , 5 , 1 , 4 , $2 + a$ equals ..... (a) 1 (b) 2 (c) 3 (d) 15
(18)	If the arithmetic mean of the values : 9 , 6 , 5 , 14 , $k$ is 7, then $k =$ ..... (a) 1 (b) 5 (c) 34 (d) 35
(19)	The mean of the values : $2 - a$ , 4 , 1 , 5 , $3 + a$ is ..... (a) 1 (b) 2 (c) 3 (d) 15
(20)	The order of the median of the set of values : 8 , 4 , 7 , 6 , 5 is ..... (a) 7 (b) 6 (c) 3 (d) 5
(21)	The order of the median of the set of values : 4 , 5 , 6 , 7 and 8 is ..... (a) third. (b) fourth. (c) fifth. (d) sixth.
(22)	If the order of the median of a set of values is the fourth , then the number of these values is ..... (a) 3 (b) 5 (c) 7 (d) 9
(23)	If the median of the set of the values : 27 , 45 , 19 , 24 and 28 is $X$ , then $X =$ ..... (a) 24 (b) 27 (c) 28 (d) 45
(24)	The median of the values : 1 , 2 , 5 , 3 and 4 is ..... (a) 3 (b) 4 (c) 5 (d) 2
(25)	The median of the values : 2 , 9 , 3 , 7 , 5 is ..... (a) 5 (b) 6 (c) 7 (d) 8

(26)	The median of the values : 3 , 7 , 5 , 8 , 2 is .....	(a) 3	(b) 5	(c) 8	(d) 7
(27)	The median of the values : 7 , 2 , 3 , 5 , 4 is .....	(a) 3	(b) 4	(c) 5	(d) 7
(28)	The median for the values 3 , 9 , 7 , 4 and 5 is .....	(a) 5	(b) 4	(c) 7	(d) 9
(29)	The median of the set of the values : 3 , 6 , 6 , 7 , 9 , 11 , 13 , 14 , 15 and 20 is .....	(a) 9	(b) 10	(c) 11	(d) 20
(30)	The median of values : 4 , 8 , 3 , 5 , 7 , 9 is .....	(a) 5	(b) 6	(c) 7	(d) 8
(31)	The median of the set of the values : 15 , 22 , 9 , 11 and 33 is .....	(a) 9	(b) 15	(c) 18	(d) 90
(32)	The median of the values : 10 , 9 , 11 , 19 , 12 is .....	(a) 9	(b) 10	(c) 11	(d) 19
(33)	The median of the set of the values : 15 , 22 , 9 , 11 and 33 is .....	(a) 9	(b) 15	(c) 18	(d) 90
(34)	The median of the values : 34 , 23 , 25 , 40 , 22 , 14 is .....	(a) 22	(b) 33	(c) 24	(d) 25
(35)	The median of the values : 41 , 23 , 15 , 30 , 20 is .....	(a) 23	(b) 15	(c) 30	(d) 20
(36)	The mode of the values : 3 , 5 , 3 , 6 , 3 and 8 is .....	(a) 3	(b) 5	(c) 6	(d) 8
(37)	The mode of the sets of values : 14 , 11 , 10 , 11 , 14 , 15 , 11 is .....	(a) 14	(b) 11	(c) 15	(d) 10
(38)	If the mode of the set of the values : 4 , 11 , 8 , 2 $X$ is 4 , then $X$ = .....	(a) 2	(b) 4	(c) 6	(d) 8
(39)	The mode of the values : 15 , 9 , $X + 1$ , 9 , 15 is 9 , then $X$ = .....	(a) 9	(b) 14	(c) 10	(d) 8
(40)	The mode of 7 , 8 , 9 , $X + 2$ and 6 is 9 then $X$ = .....	(a) 4	(b) 5	(c) 6	(d) 7

(41)

The mode of : 5 , 6 , 7 ,  $X + 2$  and 8 is 7 , then  $X = \dots\dots\dots$

- (a) 7 (b) 6 (c) 4 (d) 5

(42)

If the mode of the set of values : 4 , 11 ,  $X + 3$  , 6 is 6 , then  $X = \dots\dots\dots$

- (a) 2 (b) 3 (c) 4 (d) 6

(43)

The mode of the set of values : 5 , 9 , 5 ,  $X - 2$  , 9 is 9 , then  $X = \dots\dots\dots$

- (a) 5 (b) 57 (c) 9 (d) 11

## [2] Essay problems:

- 1 Find the arithmetic mean of the following frequency distribution :

Sets	5 –	15 –	25 –	35 –	45 –	Total
Frequency	3	10	12	10	5	40

Form the vertical table :

Set	Centre of the set « $X$ »	Frequency « $f$ »	$X \times f$
Total			

$$\text{The mean} = \frac{\text{The sum of } (X \times f)}{\text{The sum of } f} =$$



**2** Find the mean of the following data :

Sets	8 –	12 –	16 –	20 –	24 –	Total
Frequency	4	10	16	12	8	50

Form the vertical table :

Set	Centre of the set « $X$ »	Frequency « $f$ »	$X \times f$
Total			

$$\text{The mean} = \frac{\text{The sum of } (X \times f)}{\text{The sum of } f} =$$

**3** The following table shows frequency distribution of marks of 32 students in an exam :

Sets	10 –	20 –	30 –	40 –	50 –	Total
Frequency	3	6	10	8	5	32

Find the mean of this distribution.

Form the vertical table :

Set	Centre of the set « $X$ »	Frequency « $f$ »	$X \times f$
Total			

$$\text{The mean} = \frac{\text{The sum of } (X \times f)}{\text{The sum of } f} =$$

4

The following table shows the frequency distribution , find the arithmetic mean :

Sets	10 –	20 –	30 –	40 –	50 –
Frequency	10	20	25	30	15

Form the vertical table :

Set	Centre of the set « $X$ »	Frequency « $f$ »	$X \times f$
Total			

$$\text{The mean} = \frac{\text{The sum of } (X \times f)}{\text{The sum of } f} =$$

[2] Complete:

1

The most common value in a set is called .....

2

The value which is the most common of a set of values is called .....

3

The mode of a set of values is .....

4

The mode of the values : 2 , 5 , 1 , 4 , 2 is .....

5

The mode of the values : 4 , 7 , 5 , 7 , 6 , 8 , 7 , 5 is .....

6

The mode of the values : 8 , 7 , 8 , 7 , 6 , 5 , 8 is .....

7


The mode of the set of values : 13 , 12 , 4 , 13 is .....

8


The mode of the set of the values : 14 , 11 , 10 , 11 , 14 , 15 , 11 is .....



9	The mode of the values : 11 , 13 , 11 , 14 , 11 , 12 is .....
10	The mode value of : 13 , 23 , 46 , 33 , 46 , 43 , 33 , 46 , 32 is .....
11	If the mode of the set of the values : 4 , 5 , a , 3 is 4 , then a = .....
12	If the mode of the values : 3 , 6 , a , 2 , 5 is 6 , than a = .....
13	If the mode of the set of the values : 4 , 5 , a and 3 is 3 , then a = .....
14	If the mode of the values : 5 , 7 and $X + 1$ is 7 , then $X =$ .....
15	The mode of the values : 14 , 8 , $X + 1$ , 8 , 14 is 8 , then $X =$ .....
16	If the mode of the values : 12 , 7 , $X + 1$ , 7 , 12 is 7 , then $X =$ .....
17	If the mode of the set of the values : 15 , 9 , $X + 1$ , 9 and 15 is 9 , then $X =$ .....
18	If the mode of the set of the values : 15 , 9 , $X + 6$ , 9 and 15 is 9 , then $X =$ .....
19	If the mode of the values : 4 , 11 , 8 , and $2^X$ is 4 , then $X =$ .....



# GEOMETRY FOR PREPARATORY TWO FIRST TERM



# Sheet (1)

## Medians of triangle


### Definition

The median of a triangle is the line segment drawn from any vertex of this triangle to the midpoint of the opposite side of this vertex.

**For example:**

**In the opposite figure :**

If D is the midpoint of  $\overline{BC}$   
 , then  $\overline{AD}$  is a median of  $\triangle ABC$



### Notice that :

Any triangle has three medians.


### Theorem 1

The medians of a triangle are concurrent.

**For example:**

**In the opposite figure :**

$\overline{AD}$  ,  $\overline{BF}$  and  $\overline{CE}$  are the three medians of  $\triangle ABC$  ,  
 and they are concurrent at M  
 (i.e.  $\overline{AD} \cap \overline{BF} \cap \overline{CE} = \{M\}$ )



### Theorem 2


The point of concurrence of the medians of the triangle divides each median in the ratio of 1 : 2 from its base.

**For example:**

**In the opposite figure :**

In  $\triangle ABC$  , M is the point of concurrence of its medians , then :

- 1  $MD = \frac{1}{2} AM$       If  $AM = 6$  cm. , then  $MD = 3$  cm.
- 2  $CM = 2 FM$       If  $FM = 4$  cm. , then  $CM = 8$  cm.





**Remark**


The point of concurrence of the medians of the triangle divides each of them in the ratio of 2 : 1 from the vertex.

**Fact**

The point which divides the median in a triangle by the ratio of 1 : 2 from the base is the point of intersection of the medians of this triangle.


**In the opposite figure :**

If  $\overline{AD}$  is a median in  $\triangle ABC$  and  $M \in \overline{AD}$  such that  $AM = 2 MD$  ,  
then M is the point of intersection of the medians of  $\triangle ABC$



**[1] Complete:**


- 1 In  $\triangle ABC$  : if the point X is the midpoint of  $\overline{BC}$  , then  $\overline{AX}$  is called .....
- 2 The medians of the triangle are .....
- 3 The medians of the triangle intersect at .....
- 4 The point of intersection of the medians of a triangle divides each median in the ratio ..... from the vertex.
- 5 The points of concurrence of the medians of the triangle divides each median in the ratio ..... : ..... from the base.
- 6 The point of intersection of the medians of the triangle divides each of them by the ratio 1 : 2 from .....
- 7 The point which divides the median of the triangle in the ratio 1 : 2 from the base is the point of .....
- 8 **In the opposite figure :**  
If M is the point of intersection of the medians of  $\triangle ABC$  , then  $AM = \dots\dots\dots AD$



9

In the opposite figure :

If :  $MF = 2$  cm., then  $DF = \dots\dots\dots$




10

In the opposite figure :

In  $\triangle ABC$  ,  $M$  is the point of concurrence of the medians

,  $MC = 8$  cm.

, then  $DM = \dots\dots\dots$  cm.



## [2] Essay problems:


1

In the opposite figure :

$E$  is the midpoint of  $\overline{AB}$  ,  $D$  is the midpoint of  $\overline{BC}$

$\overline{AD} \cap \overline{CE} = \{M\}$  ,  $MC = 5$  cm. and  $MD = 2$  cm.

**Find :** The length of each of  $\overline{AD}$  and  $\overline{ME}$ .



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


2

In the opposite figure :

$F$  ,  $N$  are midpoints of  $\overline{AB}$  ,  $\overline{AC}$  respectively ,  $\overline{BN} \cap \overline{CF} = \{M\}$  ,

if :  $AB = 8$  cm. ,  $AC = 10$  cm. ,  $BM = 4$  cm. and  $CF = 9$  cm.

**Find :** the perimeter of figure AFMN



.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

3 In the opposite figure :

ABC is a triangle in which  $\overline{CD}$  ,  
 $\overline{BE}$  two medians intersect at M ,  
 if : DC = 9 cm. , BM = 4 cm. , BC = 8 cm.

Find : The perimeter of  $\Delta MDE$




[1] Choose the correct answer:

- |   |  |
|---|--|
| 1 | The medians of the triangle intersect at ..... point.<br>(a) 1 (b) 2 (c) 3 (d) 4   |
| 2 | The right-angled triangle has ..... medians.<br>(a) 0 (b) 1 (c) 2 (d) 3  |
| 3 | The number of medians in the right-angled triangle = .....<br>(a) 3 (b) 2 (c) 1 (d) 0  |
| 4 | The point of intersection of the medians in the triangle divides each of them by the ratio ..... from the vertex.<br>(a) 1 : 3 (b) 3 : 1 (c) 2 : 1 (d) 1 : 2 |
| 5 | The point of concurrence of the medians of the triangle divides each median in the ratio of ..... from the base.<br>(a) 1 : 2 (b) 1 : 3 (c) 2 : 1 (d) 3 : 1  |




- 6 If  $\overline{AD}$  is a median of triangle ABC , and M is the point of intersection of the medians , then  $AM = \dots\dots\dots AD$   
 (a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$
- 7  $\overline{AD}$  is a median in  $\triangle ABC$  , M is the point of intersection of its medians , then  $AM = \dots\dots\dots MD$   
 (a) 2 (b)  $\frac{1}{2}$  (c) 3 (d)  $\frac{1}{3}$
- 8 If  $\overline{XE}$  is a median in  $\triangle XYZ$  , M is the point of intersection of its medians , then  $EM = \dots\dots\dots XE$   
 (a)  $\frac{1}{2}$  (b) 2 (c)  $\frac{1}{3}$  (d)  $\frac{2}{3}$
- 9 In  $\triangle ABC$  : If  $AD = 6$  cm. is a median and M is a point of concurrent , then  $MA = \dots\dots\dots$  cm.  
 (a) 6 cm. (b) 3 cm. (c) 2 cm. (d) 4 cm.
- 10 If  $\overline{AD}$  is a median of  $\triangle ABC$  , M is the point of intersection of its medians and  $AM = 6$  cm. , then  $AD = \dots\dots\dots$   
 (a) 12 cm. (b) 6 cm. (c) 18 cm. (d) 9 cm.

- 11 In the opposite figure :  
 $\overline{AD}$  is a median in  $\triangle ABC$  , M is the point of intersection of the medians ,  $MD = 2$  cm. , then  $AD = \dots\dots\dots$  cm.  
 (a) 2 (b) 4 (c) 6 (d) 8



## [2] Essay problems:

- 1 In the opposite figure :  
 ABC is a triangle , X bisects  $\overline{AB}$  , Y bisects  $\overline{BC}$   
 ,  $XY = 5$  cm. ,  $\overline{XC} \cap \overline{AY} = \{M\}$   
 where  $CM = 8$  cm. ,  $YM = 3$  cm.  
 Find with proof the length of :  $\overline{AC}$  ,  $\overline{MX}$  ,  $\overline{AM}$



.....

.....

.....

.....

.....

.....

.....

.....



2


**In the opposite figure :**

ABC is a triangle in which  $\overline{CD}$  ,

$\overline{BH}$  are medians intersect at M ,

MC = 6 cm. , BC = 8 cm. , MB = 4 cm.

**Find with proof :** The perimeter of  $\Delta MDH$



3


**In the opposite figure :**

D and E are the midpoints of  $\overline{AB}$  and  $\overline{AC}$  respectively

,  $\overline{BE} \cap \overline{CD} = \{M\}$  , If AB = 6 cm. , AC = 10 cm.

, BM = 4 cm. and CD = 9 cm.

**Find the perimeter of the figure : ADME**



## Sheet (2)

# Medians of triangle (Follow)


### Theorem 3

In the right-angled triangle, the length of the median from the vertex of the right angle equals half the length of the hypotenuse.

**For example:**

**In the opposite figure :**

$\triangle ABC$  is a right-angled triangle at B ,  
D is the midpoint of  $\overline{AC}$  and  $AC = 10$  cm. ,  
then  $DB = 5$  cm.




### The converse of theorem 3

If the length of the median drawn from a vertex of a triangle equals half the length of the opposite side to this vertex, then the angle at this vertex is right.

**For example:**

**In the opposite figure :**

If  $\overline{BD}$  is a median in  $\triangle ABC$  ,  
 $BD = 3$  cm. and  $AC = 6$  cm. ,  
then  $m(\angle ABC) = 90^\circ$  "because  $BD = \frac{1}{2} AC$  "



### Corollary

The length of the side opposite to the angle of measure  $30^\circ$  in the right-angled triangle equals half the length of the hypotenuse.


i.e.

**In the opposite figure :**

If  $\triangle ABC$  is right-angled at B and  
 $m(\angle C) = 30^\circ$  , then  $AB = \frac{1}{2} AC$


**For example:**

If  $AC = 20$  cm. , then  $AB = 10$  cm.



# [1] Complete:

- 1 In the right-angled triangle the length of the median from the vertex of the right angle equal ..... the length of the hypotenuse.
- 2 In the right-angled triangle , the length of the median from the vertex of the right angle equals .....
- 3 If the length of the median drawn from a vertex of a triangle equals half the length of the opposite side to this vertex in length , then .....
- 4 The length of the side opposite to the angle of measure  $30^\circ$  in the right-angled triangle equals ..... the length of the hypotenuse.
- 5 The length of side opposite to the angle whose measure =  $30^\circ$  in the right-angled triangle = .....
- 6 The length of the hypotenuse on the right-angled triangle equals ..... the length of a side opposite to the angle of measure  $30^\circ$
- 7 In  $\Delta LMN$  : If  $m(\angle L) = 30^\circ$  ,  $m(\angle N) = 60^\circ$  ,  $NM = 4$  cm. , then  $LN = \dots\dots\dots$  cm.
- 8 If ABC is a right-angled triangle at B ,  $AB = 6$  cm. ,  $BC = 8$  cm. , if  $\overline{BD}$  is a median of triangle ABC , then  $BD = \dots\dots\dots$  cm.
- 9 In  $\Delta ABC$  ,  $m(\angle C) = 60^\circ$  ,  $m(\angle B) = 90^\circ$  ,  $AC = 8$  cm. , then  $BC = \dots\dots\dots$  cm.
- 10 In  $\Delta ABC$  if  $m(\angle A) = 30^\circ$  and  $m(\angle B) = 90^\circ$  , then  $BC = \dots\dots\dots AC$
- 11 If ABC : Is a right-angled at B ,  $AB = \frac{1}{2} AC$  , then  $m(\angle C) = \dots\dots\dots$
- 12 If ABC is a right-angled triangle at B and  $AB = \frac{1}{2} AC$  , then  $m(\angle A) = \dots\dots\dots$
- 13 ABC is a right-angled triangle at B , if  $AC = 2 BC$  , then  $m(\angle C) = \dots\dots\dots^\circ$
- 14 In the opposite figure :  
The perimeter of  $\Delta ABD = \dots\dots\dots$  cm.





15


In the opposite figure :

D is the midpoint of  $\overline{AC}$

,  $m(\angle E) = 30^\circ$

,  $AC = 10$  cm.

Find the length of :  $\overline{BE}$



## [2] Essay problems:


1

In the opposite figure :  $\triangle ABC$  ,  $AC = 8$  cm. ,

$m(\angle BAC) = 60^\circ$  ,  $m(\angle ABC) = 90^\circ$  ,

D is the midpoint of  $\overline{AC}$

Find : The perimeter of  $\triangle ABD$




**2** In the opposite figure :

$m(\angle B) = 90^\circ$  ,  $m(\angle C) = 30^\circ$  ,  $\overline{BD}$  is a median ,  $AB = 4$  cm. ,

Complete :

$AC = \dots\dots\dots$  cm. ,  $BD = \dots\dots\dots$  cm. ,  $AD = \dots\dots\dots$  cm.




**3** In the opposite figure :

$\Delta ABC$  in which  $m(\angle B) = 90^\circ$  ,  $AC = 10$  cm. ,

$m(\angle C) = 30^\circ$  ,  $EC = EB$  ,  $AD = DC$

Find with proof : ① The perimeter of  $\Delta ABD$

② The length of  $\overline{DF}$




**4** In the opposite figure :

$m(\angle B) = 90^\circ$  ,

$m(\angle ACB) = 30^\circ$  ,

E , F are midpoints of  $\overline{AD}$  ,  $\overline{DC}$

Prove that :  $AB = EF$




5 In the opposite figure :

$$m(\angle ABC) = m(\angle ADC) = 90^\circ ,$$

$m(\angle ACB) = 30^\circ$  , and  $\overline{DE}$  is a median of  $\triangle ADC$  ,

If  $AB = 3$  cm.

**Find :** The length of  $\overline{DE}$




6 In the opposite figure :

$$m(\angle ABC) = m(\angle BDE) = 90^\circ$$

$$, m(\angle E) = 30^\circ$$

, D is the midpoint of  $\overline{AC}$

**Prove that :**  $AC = BE$





**[1] Choose the correct answer:**

- 1 The length of the hypotenous of the right-angled triangle = ..... the length of the median which drawn from the vertex of the right-angle.  
(a) half (b) twice (c) third (d) quarter
- 2 The length of the median drawn from the vertex of right angle in the right-angled triangle = ..... the length of the hypotenuse of the triangle.  
(a) 2 (b)  $\frac{1}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$
- 3 In the right-angled triangle , the length of the median from the vertex of the right angle equals ..... the length of hypotenuse.  
(a) half (b) twice (c) third (d) forth
- 4 If  $\Delta ABC$  is a right-angled at B ,  $AB = 6$  cm. ,  $BC = 8$  cm. , then the length of the medians drawn from B is ..... cm.  
(a) 10 (b) 8 (c) 6 (d) 5
- 5 In  $\Delta ABC$  which is right at B , if  $AC = 20$  cm. , then the length of the median of the triangle drawn from B equals .....  
(a) 10 cm. (b) 8 cm. (c) 6 cm. (d) 5 cm.
- 6 In  $\Delta ABC$  ,  $m(\angle B) = 90^\circ$  ,  $AC = 12$  cm. and  $\overline{BD}$  is a median in  $\Delta ABC$  , then  $BD =$  ..... cm.  
(a) 12 (b) 6 (c) 24 (d) 10
- 7 The length of the side opposite to the angle of measure  $30^\circ$  in the right-angled ..... the length of the hypotenuse.  
(a) twice (b) half (c) square (d) equals
- 8 Triangle ABC : If  $m(\angle A) = 30^\circ$  ,  $m(\angle B) = 90^\circ$  , then  $BC =$  .....  
(a)  $\frac{1}{2} AB$  (b)  $\frac{1}{2} AC$  (c)  $2 AB$  (d)  $2 AC$
- 9 In  $\Delta ABC$  if :  $m(\angle B) = 90^\circ$  and  $m(\angle A) = 60^\circ$  , then  $AC =$  .....  $AB$   
(a) 2 (b) = (c)  $\frac{1}{2}$  (d)  $\frac{1}{3}$
- 10  $\Delta ABC$  : if  $m(\angle A) = 30^\circ$  and  $m(\angle B) = 90^\circ$  , then  $AC =$  .....  
(a)  $\frac{1}{2} BC$  (b)  $2 BC$  (c)  $2 AB$  (d)  $BC$




- |    |   |
|----|---|
| 11 | In $\Delta ABC$ : $m(\angle A) = 30^\circ$ , $m(\angle B) = 90^\circ$ , $AC = 10$ cm. , then $BC = \dots\dots\dots$ cm.<br>(a) 20 (b) 15 (c) 10 (d) 5     |
| 12 | In $\Delta XYZ$ , if $m(\angle Y) = 90^\circ$ , $m(\angle X) = 30^\circ$ and $XZ = 20$ cm. , then $ZY = \dots\dots\dots$ cm.<br>(a) 5 (b) 8 (c) 20 (d) 10 |
| 13 | In the rectangle $ACBD$ , if $AC = 10$ cm. , then $BD = \dots\dots\dots$<br>(a) 5 (b) 10 (c) 15 (d) 20  |

## [2] Complete:


- |   |   |
|---|---|
| 1 | In the right-angled triangle , the length of the median from the vertex of the right angle equals .....   |
| 2 | If the length of the median drawn from a vertex of a triangle equals half the length of the opposite side to this vertex in length , then .....               |
| 3 | The length of the side opposite to the angle of measure $30^\circ$ in the right-angled triangle equals ..... the length of the hypotenuse.                    |
| 4 | The length of side opposite to the angle whose measure = $30^\circ$ in the right-angled triangle = .....  |
| 5 | The length of the hypotenuse on the right-angled triangle equals ..... the length of a side opposite to the angle of measure $30^\circ$                       |
| 6 | In $\Delta LMN$ : If $m(\angle L) = 30^\circ$ , $m(\angle N) = 60^\circ$ , $NM = 4$ cm. , then $LN = \dots\dots\dots$ cm.                                     |
| 7 | If $ABC$ is a right-angled triangle at $B$ , $AB = 6$ cm. , $BC = 8$ cm. , if $\overline{BD}$ is a median of triangle $ABC$ , then $BD = \dots\dots\dots$ cm. |
| 8 | In $\Delta ABC$ , $m(\angle C) = 60^\circ$ , $m(\angle B) = 90^\circ$ , $AC = 8$ cm. , then $BC = \dots\dots\dots$ cm.  |
| 9 | In $\Delta ABC$ if $m(\angle A) = 30^\circ$ and $m(\angle B) = 90^\circ$ , then $BC = \dots\dots\dots AC$   |

### [3] Essay problems:

- 1 In the opposite figure :  
 $m(\angle B) = 90^\circ$  and  $m(\angle C) = 30^\circ$  ,  
 $AC = 10$  cm.  
**Find :** the lengths of  $\overline{AB}$  and  $\overline{BD}$



- 2 In the opposite figure :  
 $m(\angle C) = 30^\circ$   
**Prove that :**  
 $AB = BD$



**3 In the opposite figure :**


In  $\triangle ABC$  :  $m(\angle B) = 90^\circ$  ,  $m(\angle C) = 30^\circ$

, D is the midpoint of  $\overline{AC}$  , E is the midpoint of  $\overline{BC}$

,  $AC = 12$  cm.

(1) Find length of :  $\overline{BM}$

(2) Find the perimeter of :  $\triangle ABC$




**4 In the opposite figure :**

X , Y , D are the midpoints of  $\overline{EA}$  ,  $\overline{EC}$  and  $\overline{AC}$  respectively ,

$m(\angle ABC) = 90^\circ$


**Prove that :**  $BD = YX$



5 In the opposite figure :

$m(\angle B) = 90^\circ$  ,  $m(\angle ACB) = 30^\circ$  ,  
 $AB = DH$  where H is midpoint of  $\overline{AC}$

Prove that :  $m(\angle ADC) = 90^\circ$




6 In the opposite figure :

$m(\angle BAC) = m(\angle CBE) = 90^\circ$  ,

$m(\angle BEC) = 30^\circ$  ,

D and F are the midpoints of  $\overline{BC}$   
 and  $\overline{CE}$  respectively.

Prove that :  $AD = \frac{1}{2} BF$






## Sheet (3)

# The isosceles triangle

Triangles are classified according to the lengths of their sides into three types which are :


**1** Scalene triangle.



$$AB \neq BC \neq CA$$

**2** Isosceles triangle.


(two sides are congruent).



$$DE = DF$$

**3** Equilateral triangle.

(three sides are congruent).



$$XY = YZ = ZX$$

And in the following we will study the relations between the angles in the isosceles triangle and the equilateral triangle.

### The isosceles triangle theorem

#### Theorem 1

The base angles of the isosceles triangle are congruent.


**For example:**

**In the opposite figure :**

If ABC is a triangle in which :

$$AB = AC, m(\angle A) = 40^\circ,$$

$$\text{then } m(\angle B) = m(\angle C) = \frac{180^\circ - 40^\circ}{2} = 70^\circ$$



#### Remarks

- 1** Both of the base angles in the isosceles triangle are acute.
- 2** The vertex angle in the isosceles triangle may be acute, right or obtuse angle.

#### Corollary

If the triangle is equilateral, then it is equiangular where each angle measure is  $60^\circ$


**For example:**

**In the opposite figure :**

If XYZ is a triangle in which

$$XY = YZ = ZX ,$$

$$\text{then } m(\angle X) = m(\angle Y) = m(\angle Z) = 60^\circ$$



### [1] Complete:

- 1 The two base angles in an isosceles triangle are .....
- 2  $\triangle ABC$  ,  $AB = AC$  ,  $m(\angle C) = 70^\circ$  , then  $m(\angle A) = \dots\dots\dots$
- 3 In the  $\triangle ABC$  :  $AB = AC$  ,  $m(\angle A) = 70^\circ$  , then  $m(\angle C) = \dots\dots\dots^\circ$
- 4 The  $\triangle ABC$  is an isosceles and right-angled triangle if  $m(\angle B) = 90^\circ$  , then  $m(\angle A) = m(\angle C) = \dots\dots\dots^\circ$
- 5 In  $\triangle ABC$  , if  $AB = AC$  and  $m(\angle A) = 80^\circ$  , then  $m(\angle B) = m(\angle \dots\dots\dots) = \dots\dots\dots^\circ$
- 6 In  $\triangle ABC$  : if  $AB = AC$  ,  $m(\angle B) = 60^\circ$  , then the triangle is an .....
- 7 In  $\triangle ABC$  : If  $AB = AC$  and  $m(\angle A) = 2 m(\angle C)$  , then  $m(\angle B) = \dots\dots\dots^\circ$
- 8 The length of side opposite to the angle whose measure =  $30^\circ$  in the right-angled triangle = .....
- 9 The length of the hypotenuse on the right-angled triangle equals ..... the length of a side opposite to the angle of measure  $30^\circ$
- 10 In  $\triangle LMN$  : If  $m(\angle L) = 30^\circ$  ,  $m(\angle N) = 60^\circ$  ,  $NM = 4$  cm. , then  $LN = \dots\dots\dots$  cm.
- 11 If ABC is a right-angled triangle at B ,  $AB = 6$  cm. ,  $BC = 8$  cm. , if  $\overline{BD}$  is a median of triangle ABC , then  $BD = \dots\dots\dots$  cm.
- 12 In  $\triangle ABC$  ,  $m(\angle C) = 60^\circ$  ,  $m(\angle B) = 90^\circ$  ,  $AC = 8$  cm. , then  $BC = \dots\dots\dots$  cm.



## [2] Essay problems:


1 In the opposite figure :

ACBD is a quadrilateral in which :

$$AB = BC = CA = BD$$

$$, m(\angle ABD) = 24^\circ$$

Find :  $m(\angle CAD)$




2 In the opposite figure complete :

$$x = \dots\dots\dots^\circ,$$

$$y = \dots\dots\dots^\circ,$$

$$z = \dots\dots\dots^\circ$$




3 In the opposite figure :

ABC is a triangle ,

$$AC = BC, \overrightarrow{AD} \parallel \overrightarrow{BC}, m(\angle DAC) = 40^\circ$$

Find : The measure of angles in the  $\Delta ABC$




4 In the opposite figure :

In  $\triangle ABC$  ,  $AB = AC$  ,

$BX = CY$

Prove that :  $AX = AY$



[1] Choose the correct answer:

- 1 In any isosceles triangle , the type of the base angles is .....  
 (a) acute. (b) right. (c) obtuse. (d) reflex.
- 2 The base angles of the isosceles triangle are .....  
 (a) congruent. (b) alternate. (c) corresponding. (d) supplementary.
- 3 In  $\triangle ABC$  :  $AB = AC$  ,  $m(\angle B) = 50^\circ$  , then  $m(\angle A) = \dots\dots\dots^\circ$   
 (a) 65 (b) 80 (c) 50 (d) 100
- 4 If measure of one of the two base angles of the isosceles triangle equals  $40^\circ$  then the measure of the vertex angle = .....  
 (a) 40 (b) 100 (c) 80 (d) 50
- 5 An isosceles triangle , one of its base angles has measure  $50^\circ$  , then the measure of the vertex angle = .....  
 (a)  $50^\circ$  (b)  $60^\circ$  (c)  $70^\circ$  (d)  $80^\circ$

- |    |   |                |                 |                 |                 |
|----|---|----------------|-----------------|-----------------|-----------------|
| 6  | In the isosceles triangle, if the measure of one of the two base angles is $70^\circ$ , then the measure of its vertex angle is ..... | (a) $70^\circ$ | (b) $110^\circ$ | (c) $20^\circ$  | (d) $40^\circ$  |
| 7  | The measure of one angle of the two base angles of the isosceles = $75^\circ$ , then the measure of the vertex angle = .....          | (a) $50^\circ$ | (b) $75^\circ$  | (c) $30^\circ$  | (d) $105^\circ$ |
| 8  | In a triangle ABC : If $AB = AC$ and $m(\angle A) = 40^\circ$ , then $m(\angle C) = \dots\dots\dots$                                  | (a) $40^\circ$ | (b) $70^\circ$  | (c) $140^\circ$ | (d) $50^\circ$  |
| 9  | In $\triangle ABC$ , $AB = AC$ , $m(\angle A) = 50^\circ$ , then $m(\angle B) = \dots\dots\dots$                                      | (a) $50^\circ$ | (b) $65^\circ$  | (c) $130^\circ$ | (d) $100^\circ$ |
| 10 | If the measure of an angle of the isosceles triangle is $100^\circ$ , then the measure of one of the other angles = .....             | (a) $50^\circ$ | (b) $80^\circ$  | (c) $40^\circ$  | (d) $100^\circ$ |
| 11 | If the measure of an angle of the isosceles triangles is $120^\circ$ , then the measure of one of the other angles = .....            | (a) $60^\circ$ | (b) $30^\circ$  | (c) $40^\circ$  | (d) $45^\circ$  |
| 12 | The triangle whose sides lengths are 2 cm., $(X + 1)$ cm and 5 cm. becomes an isosceles triangle when $X = \dots\dots\dots$ cm.       | (a) 1          | (b) 2           | (c) 3           | (d) 4           |
| 13 | Triangle whose sides lengths are 2 cm., $(X - 2)$ cm., 5 cm. becomes isosceles triangle when $X = \dots\dots\dots$ cm.                | (a) 3          | (b) 4           | (c) 5           | (d) 7           |
| 14 | ABC is a triangle in which $AB = AC$ and $m(\angle A) = 110^\circ$ , then $m(\angle B) = \dots\dots\dots$                             | (a) $70^\circ$ | (b) $55^\circ$  | (c) $35^\circ$  | (d) $110^\circ$ |
| 15 | $\triangle XYZ$ is an isosceles triangle in which $m(\angle X) = 100^\circ$ , then $m(\angle Y) = \dots\dots\dots^\circ$              | (a) 100        | (b) 80          | (c) 60          | (d) 40          |




## [2] Essay problems:

1 In the opposite figure :

$AB = AD$  ,  $m(\angle A) = 30^\circ$  ,

$CB = BD = CD$

Find :  $m(\angle CBA)$




2 In the opposite figure :

$ABC$  is a triangle in which :  $AC = BC$  ,

$\overline{AD} \parallel \overline{BC}$  ,  $m(\angle DAC) = 30^\circ$

Find :  $m(\angle ABC)$




3

In the opposite figure :

ABC is an equilateral triangle ,

$DB = DC$  ,  $m(\angle D) = 110^\circ$

Find with proof :  $m(\angle DBC)$  and  $m(\angle DBA)$




4

In the opposite figure :

ADE is a triangle ,  $B \in \overline{DE}$  ,  $C \in \overline{DE}$

,  $BD = CE$  ,  $AB = AC$

Prove that :  $AD = AE$





## Sheet (4)

## The converse of the isosceles triangle theorem

**Theorem 2**

If two angles of a triangle are congruent , then the two sides opposite to these two angles are congruent and the triangle is isosceles.

**Remark**

The isosceles triangle in which the measure of one of its angles =  $60^\circ$  is an equilateral triangle.


**[1] Complete:**

- |          |  |
|----------|--|
| <b>1</b> | If angles of any triangle are equal in measures , then the triangle is .....   |
| <b>2</b> | If the angles of a triangle are congruent , then the triangle is .....   |
| <b>3</b> | The measure of the exterior angle of equilateral triangle = ..... $^\circ$   |
| <b>4</b> | If the measure of one of the angles of the right-angled triangle is $45^\circ$ , then the triangle is .....            |
| <b>5</b> | In an isosceles triangle , if any angle has a measure of $60^\circ$ , the triangle is .....                            |
| <b>6</b> | In $\triangle ABC$ if : $\overline{AB} \perp \overline{BC}$ and $AB = BC$ , then $m(\angle A) = \dots\dots\dots^\circ$ |

**[2] Essay problems:****1 In the opposite figure :** $\overrightarrow{AD}$  bisects  $\angle BAC$ 

$m(\angle B) = 30^\circ$

$m(\angle C) = 70^\circ$

**Prove that :**  $\triangle ADC$  is isosceles triangle.**2** ABC is a triangle in which :  $m(\angle A) = 50^\circ$  and  $m(\angle C) = 80^\circ$ **Prove that :** this triangle ABC is an isosceles triangle.


3

In the opposite figure :

$AB = AC$  ,

$\overrightarrow{BM}$  and  $\overrightarrow{CM}$  bisect the angles  $(\angle B)$  ,  $(\angle C)$

Prove that :  $MB = MC$



4


In the opposite figure :

$BD = CE$

,  $m(\angle ABC) = m(\angle ACB)$

,  $m(\angle D) = m(\angle E) = 90^\circ$

Prove that :  $m(\angle DAB) = m(\angle CAE)$




5 In the opposite figure :

$$AB = AC, \overline{DE} \parallel \overline{AB}$$

$$\text{and } \overline{AC} \parallel \overline{DO}$$

Prove that : ①  $DE = DO$       ②  $m(\angle A) = m(\angle D)$



[1] Choose the correct answer:

- 1 The measure of exterior angle of an equilateral triangle = .....  
 (a)  $30^\circ$                       (b)  $60^\circ$                       (c)  $120^\circ$                       (d)  $180^\circ$
- 2 In  $\triangle XYZ$  : if  $XY = XZ$  , then the exterior angle at the vertex Z is .....  
 (a) acute.                      (b) obtuse.                      (c) right.                      (d) reflex.
- 3 In  $\triangle ABC$  : if  $AB = AC$  and  $m(\angle A) = 60^\circ$  , if its perimeter is 18 cm. , then  $BC = \dots\dots\dots$  cm.  
 (a) 18                      (b) 6                      (c) 3                      (d) 60



## [2] Essay problems:


1 In the opposite figure :

$$AD = AC$$

$$, m(\angle DAB) = 30^\circ$$

$$, m(\angle ABD) = 40^\circ$$

**Prove that :**  $AB = CB$




2 In the opposite figure :

ABC is a triangle in which  $AB = AC$  ,  $X \in \overline{AB}$  ,

$Y \in \overline{AC}$  and  $\overline{XY} \parallel \overline{BC}$

**Prove that :** the triangle AXY is isosceles triangle.






3 In the opposite figure :

$AB = AC$  ,  $\overrightarrow{BD}$  bisects  $\angle B$  and  $\overrightarrow{CD}$  bisects  $\angle C$


Prove that :  $\triangle DBC$  is an isosceles triangle



4 In the opposite figure :

$\overrightarrow{CD}$  bisects  $\angle ACB$  ,  $\overrightarrow{DE} \parallel \overrightarrow{CB}$

Prove that :  $\triangle ECD$  is an isosceles triangle.




5 | In the opposite figure :

$$m(\angle B) = m(\angle C), AB = (2x - 1) \text{ cm.}$$

$$AC = (x + 3) \text{ cm.}$$

$$BC = (9 - x) \text{ cm.}$$

Find with proof the perimeter of  $\triangle ABC$



## Sheet (5)

### Corollaries of the Isosceles triangle theorems

#### Corollary 1

The median of an isosceles triangle from the vertex angle bisects it and is perpendicular to the base.

In the opposite figure :


ABC is a triangle in which  $AB = AC$  and

$\overline{AD}$  is a median , then :

1  $\overline{AD}$  bisects  $\angle BAC$

i.e.  $m(\angle BAD) = m(\angle CAD)$

2  $\overline{AD} \perp \overline{BC}$



#### Corollary 2

The bisector of the vertex angle of an isosceles triangle bisects the base and is perpendicular to it.

In the opposite figure :


ABC is a triangle in which  $AB = AC$  and

$\overline{AD}$  bisects  $\angle BAC$  , then :

1 D is the midpoint of  $\overline{BC}$

i.e.  $BD = CD$

2  $\overline{AD} \perp \overline{BC}$



#### Corollary 3

The straight line drawn passing through the vertex angle of an isosceles triangle perpendicular to the base bisects each of the base and the vertex angle.

In the opposite figure :


ABC is a triangle in which  $AB = AC$  and

$\overline{AD} \perp \overline{BC}$  , then :

1 D is the midpoint of  $\overline{BC}$

i.e.  $BD = CD$

2  $m(\angle BAD) = m(\angle CAD)$



**Notice that :**

The previous three corollaries can be proved using the congruence of  $\triangle ABD$  and  $\triangle ACD$


**Axis of symmetry of a line segment**

**Definition**

The straight line perpendicular to a line segment at its middle is called the axis of symmetry for that line segment , in brief it is known as the axis of a line segment.

**In the opposite figure :**

If the straight line  $L \perp \overline{AB}$  and  $C \in$  the straight line  $L$  where  $C$  is the midpoint of  $\overline{AB}$  , then the straight line  $L$  is called the axis of  $\overline{AB}$




**Property**

Any point on the axis of symmetry of a line segment is at equal distances from its terminals (end points).

**In the opposite figure :**

If the straight line  $L$  is the axis of  $\overline{AB}$  ,  
 $D \in L$  ,  $E \in L$  and  $F \in L$  , then  
 $DA = DB$  ,  $EA = EB$  and  $FA = FB$




**The converse of the previous property is true**

i.e. If a point is at equal distances from the two terminals of a line segment , then this point lies on the axis of this line segment.

**In the opposite figure :**

If  $C$  is a point such  
 that  $CA = CB$  , then  
 the point  $C$  lies on the axis of  $\overline{AB}$



**Axis of symmetry of the isosceles triangle**


The isosceles triangle has one axis of symmetry.

It is the straight line drawn from the vertex angle perpendicular to its base.



**For example:**

If ABC is an isosceles triangle where  $AB = AC$  and  $\overrightarrow{AD} \perp \overrightarrow{BC}$ , then  $\overrightarrow{AD}$  is called the axis of symmetry of the isosceles triangle ABC




**Remarks**

- 1 The equilateral triangle has three axes of symmetry, they are the three perpendiculars drawn from its vertices to the opposite sides.

**In the opposite figure :**

The straight lines  $L_1$ ,  $L_2$  and  $L_3$  are the axes of symmetry of the equilateral triangle ABC



- 2 The scalene triangle has no axes of symmetry.

**[1] Complete:**


- 1 The ray drawn from the vertex of the isosceles triangle passing through the midpoint of the base is .....
- 2 The median of an isosceles triangle drawn from the vertex bisects ..... and is perpendicular to .....
- 3 The bisector of the vertex angle of an isosceles triangle ..... and .....
- 4 In  $\triangle XYZ$  : If  $XY = XZ$ ,  $\overrightarrow{XL} \perp \overrightarrow{YZ}$ , then  $\overrightarrow{XL}$  bisects each of ..... and .....
- 5 The straight line perpendicular to the midpoint of a line segment is called .....
- 6 In the isosceles triangle if the measure of any angle is  $60^\circ$ , then the number of axis of symmetry .....
- 7 The number of axes of symmetry of the isosceles triangle equal .....
- 8 The number of symmetrical line in an scalene triangle = .....
- 9 The number of the axes of symmetry in an equilateral triangle = .....



- 10** The number of axes of symmetry of the triangle in which the measures of two angles are  $50^\circ$  ,  $70^\circ$  = .....
- 11** In  $\triangle ABC$  : If  $AB = AC$  , then the point A lies on the axis of symmetry of .....
- 12** If D is the midpoint of  $\overline{AB}$  and  $\overrightarrow{CD} \perp \overline{AB}$  , then  $CA =$  .....
- 13** The axis of symmetry of the line segment is the straight line which .....
- 14** Any point on the axis symmetry of a line segment is at two equal distance from .....
- 15** If the point  $A \in$  the axis of symmetry of  $\overline{BC}$  , then  $AB =$  .....
- 16** The axis of symmetry of isosceles triangle is .....

## [2] Essay problems:

- 1** In the opposite figure :  
 In  $\triangle ABC$  ,  $AB = AC$  ,  
 $\overline{AD} \perp \overline{BC}$  ,  
 $AB = 13$  cm. and  $BD = 5$  cm.  
**Find :** ① The length of  $\overline{BC}$   
 ② The area of  $\triangle ABC$




2

**In the opposite figure :**

ABC is a triangle in which :  $AB = AC$  ,  $\overline{AD} \perp \overline{BC}$

$m(\angle BAC) = 100^\circ$  and  $BD = 3$  cm.

**Find :** (1)  $m(\angle BAD)$       (2) The length of  $\overline{CB}$



3


**In the opposite figure :**

$XL = XY$  ,  $ZL = ZY$  ,

M is the midpoint of  $\overline{LY}$

**Prove that :**

X , M , Z are on the same straight line.



4

**In the opposite figure :**


$\overrightarrow{AD}$  bisects  $\angle BAC$  ,

$AB = AC = 17$  cm. ,

and  $BC = 16$  cm.

**Prove that :**  $m(\angle ADB) = 90^\circ$  ,

then find the length of :  $\overline{AD}$  and the area of  $\triangle ABC$



**[1] Choose the correct answer:**

1

The axis of symmetry of a line segment is the straight line which is .....

(a) Perpendicular to it.

(b) its bisector.

(c) parallel to it.

(d) the perpendicular bisector.

2

If  $A \in$  the axis of symmetry of  $\overline{BC}$  , then  $\overline{AB} \dots\dots \overline{AC}$

(a)  $\perp$

(b)  $\equiv$

(c)  $//$

(d) =

3

If A lies on the axis of symmetry of  $\overline{XY}$  then  $AX \dots\dots AY$

(a)  $//$

(b)  $\perp$

(c) =

(d)  $\neq$

4

The number of axis of symmetry in the scalene triangle is .....

(a) 1

(b) zero

(c) 3

(d) 4

5

The number of axes of symmetry in the isosceles triangle is .....

(a) 1

(b) 2

(c) 3

(d) zero

- 6** The isosceles triangle has ..... axis (axes) of symmetry.  
(a) no (b) two (c) only one (d) three
- 7** The number of axes of symmetry in the equilateral triangle is .....  
(a) 0 (b) 2 (c) 3 (d) 1
- 8** The equilateral triangle has ..... axes of symmetry.  
(a) one (b) two (c) three (d) otherwise
- 9** The triangle which has no axes of symmetry is ..... triangles.  
(a) scalene (b) isosceles (c) equilateral (d) otherwise
- 10** If  $\Delta ABC$  has one axes of symmetry and  $m(\angle ABC) = 140^\circ$ , then  $m(\angle A) = \dots\dots\dots$   
(a)  $30^\circ$  (b)  $20^\circ$  (c)  $40^\circ$  (d)  $60^\circ$
- 11** The triangle which has three axes of symmetry is ..... triangle.  
(a) scalene (b) isosceles (c) right-angled (d) equilateral
- 12**  $\Delta ABC$  in which  $m(\angle A) = m(\angle B) = 65^\circ$ , then it has ..... axis (axes) of symmetry.  
(a) 1 (b) 2 (c) 3 (d) zero
- 13** In  $\Delta ABC$  if :  $m(\angle A) = 40^\circ$  and  $m(\angle B) = 70^\circ$ , then  $\Delta ABC$  has ..... axis (axes) of symmetry.  
(a) 3 (b) 1 (c) 2 (d) zero
- 14** The quadrilateral ABCD in which  $\overline{BD}$  is an axis of symmetry of  $\overline{AC}$  may be .....  
(a) a rhombus (b) a rectangle (c) a parallelogram (d) a trapezium
- 15**  $\Delta ABC$ ,  $AB = AC$ , D is the midpoint of  $\overline{BC}$ , then  $\overline{AD}$  is .....  
(a) median. (b) altitude.  
(c) bisector of the vertex angle. (d) all the previous.

## [2] Essay problems:

### 1 In the opposite figure :


In  $\triangle ABC$  ,  $AB = AC$  ,

$\overline{AD} \perp \overline{BC}$  ,

$AB = 13$  cm. and  $BD = 5$  cm.

Find : ① The length of  $\overline{BC}$

② The area of  $\triangle ABC$




### 2 In the opposite figure :

$ABC$  is a triangle in which :  $AB = AC$  ,  $\overline{AD} \perp \overline{BC}$

$m(\angle BAC) = 100^\circ$  and  $BD = 3$  cm.

Find : ①  $m(\angle BAD)$       ② The length of  $\overline{CB}$





**3** In the opposite figure :


In  $\triangle ABC$  :

$AB = AC$  ,  $\overrightarrow{AD}$  bisects  $\angle BAC$

and  $BD = 3$  cm.

**Prove that :**  $\overline{AD} \perp \overline{BC}$

**, then find the length of :  $\overline{CB}$**




**4** In the opposite figure :

$XL = XY$  ,  $ZL = ZY$  ,

M is the midpoint of  $\overline{LY}$

**Prove that :**

$X$  ,  $M$  ,  $Z$  are on the same straight line.



## Sheet (6)

### Comparing the measure of angles in a triangle

#### Axioms of inequality relation

For any four numbers  $a$ ,  $b$ ,  $c$  and  $d$  :

- |  |  |
|--|--|
| <p><b>1</b> If <math>a &gt; b</math>, then <math>a + c &gt; b + c</math></p> <p><b>3</b> If <math>a &gt; b</math>, <math>c &gt; 0</math>, then <math>ac &gt; bc</math></p> <p><b>5</b> If <math>a &gt; b</math>, <math>c &gt; d</math>, then <math>a + c &gt; b + d</math></p> | <p><b>2</b> If <math>a &gt; b</math>, then <math>a - c &gt; b - c</math></p> <p><b>4</b> If <math>a &gt; b</math>, <math>b &gt; c</math>, then <math>a &gt; c</math></p> |
|--|--|

#### Remember that :

The measure of any exterior angle of a triangle is greater than the measure of any interior angle of the triangle except its adjacent angle.

#### Theorem


In a triangle, if two sides have unequal lengths, then the longer is opposite to the angle of the greater measure.

#### Remark

The greatest angle in measure of the triangle is opposite to the longest side of the triangle and its measure is greater than  $60^\circ$ , and the smallest angle in measure of the triangle is opposite to the shortest side of the triangle and its measure is less than  $60^\circ$


i.e. In  $\triangle ABC$  :

If  $AB > BC > AC$ , then  $m(\angle C) > m(\angle A) > m(\angle B)$   
 $, m(\angle C) > 60^\circ$  and  $m(\angle B) < 60^\circ$



# [1] Complete:


- 1 The length of two sides in the triangle are not equal , then the greatest side in length is opposite to .....
- 2 In a triangle , if two sides have unequal lengths , the longer is opposite to the angle of the .....
- 3 In triangle ABC , if  $BC > AB$  , then  $m(\angle A)$  .....  $m(\angle C)$
- 4 In  $\triangle ABC$  :  $AB > AC$  , then  $m(\angle C)$  .....  $m(\angle B)$
- 5 In  $\triangle ABC$  , if  $AB > BC > AC$  , then the smallest angle in measure of it is angle .....
- 6 In  $\triangle ABC$  : if the point X is the midpoint of  $\overline{BC}$  , then  $\overline{AX}$  is called .....
- 7 The medians of the triangle are .....
- 8 The medians of the triangle intersect at .....
- 9 The point of intersection of the medians of a triangle divides each median in the ratio ..... from the vertex.
- 10 The points of concurrence of the medians of the triangle divides each median in the ratio ..... : ..... from the base.
- 11 The point of intersection of the medians of the triangle divides each of them by the ratio 1 : 2 from .....
- 12 The point which divides the median of the triangle in the ratio 1 : 2 from the base is the point of .....
- 13 **In the opposite figure :**  
If M is intersection point of medians  
and  $m(\angle B) = 90^\circ$  ,  $MF = 1.5$  cm.  
 , then the length of  $\overline{AC} =$  .....



## [2] Essay problems:

1 In the opposite figure :


Arrange the angles of  $\triangle ABC$   
descendingly due to their measures



2 In the opposite figure :

ABCD is a quadrilateral in which :  $AD = DC$  ,  
 $BC > AB$

Prove that :  $m(\angle BAD) > m(\angle BCD)$






**3** In the opposite figure :

$AD > DC$

and  $AB > BC$

**Prove that :**  $m(\angle BCD) > m(\angle BAD)$




**[1] Choose the correct answer:**

- 1 In  $\triangle ABC$  ,  $AB > AC$  , then  $m(\angle C) \dots\dots\dots m(\angle B)$   
 (a)  $<$  (b)  $>$  (c)  $=$  (d)  $\leq$
- 2 In  $\triangle ABC$  if  $AB > AC$  , then  $m(\angle B) \dots\dots\dots m(\angle C)$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\geq$
- 3 In  $\triangle ABC$  ,  $AB > AC$  ,  $m(\angle C) = 70^\circ$  , then  $m(\angle B)$  may be .....  
 (a)  $70^\circ$  (b)  $50^\circ$  (c)  $80^\circ$  (d)  $75^\circ$
- 4 In  $\triangle ABC$  : If  $BC > AB$  , then  $m(\angle A) \dots\dots\dots m(\angle C)$   
 (a)  $=$  (b)  $<$  (c)  $\leq$  (d)  $>$
- 5 In the triangle XYZ , if  $XY > ZX$  , then  $m(\angle Y) \dots\dots\dots m(\angle Z)$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\geq$
- 6 In  $\triangle ABC$  :  $AB = AC$  ,  $m(\angle B) = 65^\circ$  , then :  $AC \dots\dots\dots BC$   
 (a)  $<$  (b)  $>$  (c)  $=$  (d)  $\leq$



- 7 In  $\triangle ABC$  : If  $AB = 9$  cm. ,  $BC = 6$  cm. ,  $AC = 7$  cm. , then the smallest angle is .....  
 (a)  $\angle BAC$  (b)  $\angle ABC$  (c)  $\angle ACB$  (d)  $\angle BCA$
- 8 The medians of the triangle intersect at ..... point.  
 (a) 1 (b) 2 (c) 3 (d) 4
- 9 The right-angled triangle has ..... medians.  
 (a) 0 (b) 1 (c) 2 (d) 3
- 10 The point of concurrence of the medians of the triangle divides each median in the ratio of ..... from the base.  
 (a) 1 : 2 (b) 1 : 3 (c) 2 : 1 (d) 3 : 1
- 11 If  $\overline{AD}$  is a median of triangle  $ABC$  , and  $M$  is the point of intersection of the medians , then  $AM = \dots\dots\dots AD$   
 (a)  $\frac{1}{3}$  (b)  $\frac{2}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{1}{4}$
- 12  $\overline{AD}$  is a median in  $\triangle ABC$  ,  $M$  is the point of intersection of its medians , then  $AM = \dots\dots\dots MD$   
 (a) 2 (b)  $\frac{1}{2}$  (c) 3 (d)  $\frac{1}{3}$
- 13 If  $\overline{XE}$  is a median in  $\triangle XYZ$  ,  $M$  is the point of intersection of its medians , then  $EM = \dots\dots\dots XE$   
 (a)  $\frac{1}{2}$  (b) 2 (c)  $\frac{1}{3}$  (d)  $\frac{2}{3}$
- 14 In the opposite figure :  
 $\overline{AD}$  is a median in  $\triangle ABC$  ,  $M$  is the point of intersection of the medians ,  $MD = 2$  cm. , then  $AD = \dots\dots\dots$  cm.  
 (a) 2 (b) 4 (c) 6 (d) 8



**[2] Essay problems:**

- 1 In  $\triangle ABC$  if :  $AB = 14$  cm. ,  $BC = 6$  cm. and  $AC = 10$  cm. Arrange the angles of  $\triangle ABC$  ascendingly due to their measures.

.....

.....

.....

.....

.....

.....

- 2  $\triangle XYZ$  in which ,  $XY = 8$  cm. ,  $YZ = 10$  cm. and  $ZX = 7$  cm. ordered the measures of its angles ascending.

.....

.....


.....

.....

.....

.....

- 3 In the opposite figure :  
 $ABC$  is a triangle ,  
 $AB > AC$  ,  $\overline{XY} \parallel \overline{BC}$   
 Prove that :  $m(\angle AYX) > m(\angle AXY)$



.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

4 In the opposite figure :

$\overline{ED} \parallel \overline{BC}$  ,

$AC > AB$

Prove that :  $AE > AD$




5 In the opposite figure :

$AB = 8 \text{ cm.}$  ,

$BC = 7 \text{ cm.}$  ,

$CD = 4 \text{ cm.}$  ,  $AD = 6 \text{ cm.}$

Prove that :  $m(\angle BCD) > m(\angle BAD)$




**6** In the opposite figure :

XYZL is a quadrilateral ,

$$XL = LZ, YZ > XY$$

**Prove that :**  $m(\angle LXY) > m(\angle LZY)$





## Sheet (7)

### Comparing the lengths of sides in a triangle

#### Theorem

In a triangle, if two angles are unequal in measure, then the greater angle in measure is opposite to a side greater in length than that opposite to the other angle.


#### Corollaries

##### Corollary 1

In the right-angled triangle, the hypotenuse is the longest side.

**In the opposite figure :**

If  $\triangle ABC$  is right-angled at  $B$ , then  $m(\angle B) > m(\angle A)$ ,  
 $m(\angle B) > m(\angle C)$  because  $\angle B$  is a right angle and each of  
 $\angle A$  and  $\angle C$  is acute, so we find that :  
 $AC > BC$  and  $AC > AB$  (according to the previous theorem).



**Notice that :**


In the obtuse-angled triangle, the side opposite to the obtuse angle is the longest side in the triangle.

##### Corollary 2

The length of the perpendicular line segment drawn from a point outside a straight line to this line is shorter than any line segment drawn from this point to the given straight line.

**In the opposite figure :**

If  $C \notin \overleftrightarrow{AB}$  and  $D \in \overleftrightarrow{AB}$  such that  $\overline{CD} \perp \overleftrightarrow{AB}$ ,  
 then  $\overline{CB}$  is the hypotenuse in  $\triangle CBD$   
 which is right-angled at  $D$ ,



$\overline{CA}$  is the hypotenuse in  $\triangle CDA$  which is right-angled at  $D$  and so on ...

According to corollary 1, we find that  $CB > CD$ ,  $CA > CD$  and so on ...

i.e.  $CD < CB$  and  $CD < CA$

#### Definition

The distance between any point and a given straight line is the length of the perpendicular line segment drawn from this point to the given line.



**[1] Complete:**

- 1 In the right-angled triangle the longest side in it called .....
- 2 The longest side length in the right-angled triangle is .....
- 3 If XYZ is a right-angled triangle at Y , then the longest side is .....
- 4 In a triangle if two angles have unequal in measure , then the greatest angle in measure is .....
- 5 The smallest angle of a triangle (in measure) is opposite to .....
- 6 In any triangle the greatest angle in measure is opposite to .....
- 7 If ABC is an obtuse-angled triangle at C , then AB ..... BC
- 8 If :  $x > y$  , z is positive number then :  $xz > \dots\dots\dots$
- 9  $\Delta ABC$  in which :  $m(\angle A) = 100^\circ$  , then the greatest side in length is .....
- 10 The longest side in the triangle ABC in which  $m(\angle B) = 105^\circ$  is .....
- 11  $\Delta ABC$  in which  $m(\angle A) = 110^\circ$  , then the greatest side in length is .....
- 12  $\Delta ABC$  in which :  $m(\angle C) = 112^\circ$  , then the longest side is .....
- 13 In  $\Delta ABC$  : If  $m(\angle B) = 120^\circ$  , then the longest side in  $\Delta ABC$  is .....
- 14 In  $\Delta DEF$  if  $m(\angle E) = 125^\circ$  , then the longest side in this triangle is .....
- 15 In  $\Delta ABC$  : If  $m(\angle A) = 130^\circ$  , then the longest side is .....
- 16 In triangle ABC , if  $m(\angle A) = 70^\circ$  ,  $m(\angle B) = 30^\circ$  , then the longest side in length is .....
- 17 In  $\Delta ABC$  : if  $m(\angle A) = 67^\circ$  and  $m(\angle B) = 33^\circ$  , then  $AB > \dots\dots\dots > \dots\dots\dots$
- 18  $\Delta ABC$  in which  $m(\angle B) = 70^\circ$  and  $m(\angle C) = 35^\circ$  the longest side in length is .....

19 In  $\triangle ABC$  ,  $m(\angle A) = 50^\circ$  ,  $m(\angle B) = 65^\circ$  , then the number of axes of symmetry equals .....


20 In  $\triangle ABC$  if :  $m(\angle A) = 50^\circ$  and  $m(\angle B) = 60^\circ$  , then the longest side in this triangle is .....

21 In  $\triangle ABC$   $m(\angle B) = 70^\circ$  and  $m(\angle C) = 60^\circ$  then  $AC$  .....  $AB$

22 In the isosceles triangle if :  $AB = AC$  ,  $m(\angle A) = 70^\circ$  , then  $AB < \dots\dots\dots$

23 In the triangle  $ABC$  : if  $m(\angle B) - m(\angle A) > m(\angle C)$  , then  $AC$  .....  $AB$

## [2] Essay problems:

1   $ABC$  is a triangle in which :  $m(\angle A) = 40^\circ$  and  $m(\angle B) = 75^\circ$   
Order the lengths of the sides of the triangle descendingly.

.....


.....


.....

.....

.....

.....

2  In the opposite figure :  
 $ABC$  is an obtuse-angled triangle at  $B$   
,  $\overline{DE} \parallel \overline{BC}$   
Prove that :  
 $AE > AD$



.....

.....

.....

.....

.....

.....

.....

.....

.....


 ABC is a right-angled triangle at B ,  $D \in \overline{AC}$  and  $E \in \overline{BC}$  where  $AD = BE$

**Prove that :  $m(\angle CED) > m(\angle CDE)$**

 **In the opposite figure :**

$$\overline{AB} \cap \overline{CD} = \{M\}, \overline{AC} \perp \overline{CD} \text{ and } \overline{BD} \perp \overline{CD}$$

**Prove that :**

$$AB > CD$$




[1] Choose the correct answer:


- 1  $\Delta XYZ$  ,  $m(\angle X) = 60^\circ$  ,  $m(\angle Y) = 40^\circ$  , then  $XZ$  .....  $XY$   
 (a)  $<$  (b)  $>$  (c)  $=$  (d) nothing.
- 2 ABC is a triangle in which :  $m(\angle B) = 70^\circ$  ,  $m(\angle C) = 50^\circ$  , then  $AC$  .....  $AB$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\equiv$
- 3 In a triangle ABC :  $m(\angle B) = 75^\circ$  ,  $m(\angle C) = 50^\circ$  , then  $BC$  .....  $AB$   
 (a)  $<$  (b)  $>$  (c)  $=$  (d)  $\equiv$
- 4 ABC is a triangle in which :  $m(\angle B) = 80^\circ$  ,  $m(\angle C) = 50^\circ$  , then  $BC$  .....  $AB$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\equiv$
- 5 If :  $m(\angle A) = 50^\circ$  and  $m(\angle B) = 60^\circ$  in triangle ABC then  $AB$  .....  $AC$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\leq$
- 6 In a triangle ABC : If  $m(\angle A) = 80^\circ$  ,  $m(\angle C) = 60^\circ$  , then  $AB$  .....  $BC$   
 (a)  $>$  (b)  $<$  (c)  $=$  (d)  $\geq$
- 7 Triangle ABC : If  $m(\angle B) = 70^\circ$  ,  $m(\angle C) = 60^\circ$  , then  $BC$  .....  $AB$   
 (a)  $<$  (b)  $>$  (c)  $=$  (d)  $\geq$
- 8 In  $\Delta LMN$  , if  $m(\angle N) = 75^\circ$  ,  $m(\angle M) = 60^\circ$  , then  $LM$  .....  $LN$ .  
 (a)  $>$  (b)  $<$  (c)  $=$  (d) twice
- 9  $\Delta ABC$  ,  $m(\angle B) = 90^\circ$  , then  $AB$  .....  $AC$   
 (a)  $>$  (b)  $=$  (c)  $<$  (d)  $\geq$
- 10 In  $\Delta XYZ$  : If  $m(\angle X) = 30^\circ$  and  $m(\angle Y) = 80^\circ$  , then .....  
 (a)  $XY < XZ$  (b)  $XY > XZ$  (c)  $XY = XZ$  (d)  $XY < YZ$
- 11 In  $\Delta ABC$  :  $m(\angle A) = 60^\circ$  and  $m(\angle C) = 45^\circ$  , then .....  
 (a)  $AB < AC$  (b)  $AB = AC$  (c)  $AB > AC$  (d)  $\overline{AB} \equiv \overline{AC}$

- 12** KLM is a triangle in which  $m(\angle K) = 50^\circ$  and  $m(\angle M) = 60^\circ$ , then which of the following statement is true ? .....
- (a)  $KL = KM$       (b)  $KM > KL$       (c)  $KM < ML$       (d)  $LM > KL$
- 13** The triangle in which the measure of two angles are  $74^\circ$  and  $53^\circ$  is ..... triangle.
- (a) a right-angled      (b) an isosceles      (c) an equilateral      (d) a scalene
- 14** In  $\Delta ABC$  : If  $AB > AC$ ,  $m(\angle C) = 70^\circ$ , then  $m(\angle B)$  may equal .....
- (a)  $70^\circ$       (b)  $50^\circ$       (c)  $80^\circ$       (d)  $75^\circ$
- 15** In  $\Delta ABC$ , if  $m(\angle A) = 50^\circ$  and  $m(\angle B) = 30^\circ$ , then the shortest side in the triangle ABC is .....
- (a)  $\overline{AB}$       (b)  $\overline{CB}$       (c)  $\overline{AC}$       (d)  $\overline{BC}$
- 16**  $\Delta ABC$  which :  $m(\angle A) = 50^\circ$ ,  $m(\angle B) = 60^\circ$  the longest side of it is .....
- (a)  $\overline{AB}$       (b)  $\overline{AC}$       (c)  $\overline{BC}$       (d)  $\overline{CB}$
- 17** In  $\Delta ABC$  if :  $m(\angle B) = 60^\circ$  and  $m(\angle C) = 50^\circ$ , then the shortest side in triangle ABC is .....
- (a)  $\overline{AC}$       (b)  $\overline{BC}$       (c)  $\overline{BC}$       (d)  $\overline{AB}$
- 18** In the triangle ABC, if  $m(\angle B) = 90^\circ$ , then the greatest side in length is .....
- (a)  $\overline{AB}$       (b)  $\overline{BC}$       (c)  $\overline{AC}$       (d)  $\overline{XY}$
- 19** In  $\Delta ABC$  if :  $m(\angle B) = 130^\circ$ , then the longest side of it is .....
- (a)  $\overline{BC}$       (b)  $\overline{AC}$       (c)  $\overline{AB}$       (d) it's median
- 20** In the triangle ABC : If  $m(\angle B) > m(\angle C)$ , then  $AB$  .....  $AC$
- (a)  $<$       (b)  $>$       (c)  $=$       (d) otherwise
- 21** In  $\Delta ABC$  : if  $m(\angle B) > m(\angle C)$ , then  $AC$  .....  $AB$
- (a)  $>$       (b)  $<$       (c)  $=$       (d)  $\leq$
- 22** In  $\Delta ABC$ , if  $m(\angle B) > m(\angle C)$ , then .....
- (a)  $AB < AC$       (b)  $AB = AC$       (c)  $AB > AC$       (d)  $\overline{AB} \equiv \overline{AC}$




- |           |   |
|-----------|---|
| <b>23</b> | In $\triangle ABC$ : $m(\angle A) < m(\angle C) < m(\angle B)$ , then .....<br>(a) $AB > AC$ (b) $BC > AC$ (c) $AC > AB$ (d) $BC > AB$  |
| <b>24</b> | The triangle ABC is obtuse-angled triangle at B , then the longest side is .....<br>(a) AB                                      (b) BC                                      (c) AC                                      (d) AD                              |
| <b>25</b> | $\triangle XYZ$ is right-angled at Y , then $XZ$ ..... $YZ$<br>(a) =                                      (b) >                                      (c) $\leq$ (d) <   |
| <b>26</b> | In $\triangle ABC$ : $m(\angle B) + m(\angle C) = 3 m(\angle A)$ , then $m(\angle A) = \dots\dots\dots^\circ$<br>(a) 30                                      (b) 60                                      (c) 45                                      (d) 90 |

## [2] Essay problems:

- 1**  **In the opposite figure :**  
 ABC is a triangle ,  $D \in \overrightarrow{CB}$  ,  
 $E \in \overrightarrow{AC}$  ,  $m(\angle ABD) = 110^\circ$   
 and  $m(\angle BCE) = 120^\circ$

**Prove that :  $AB > BC$**



2




In the opposite figure :

$$AB = AC, m(\angle ABC) = 65^\circ$$

$$, m(\angle ACD) = 20^\circ, A \in \overline{BD}$$

Prove that :  $AB > AD$



3




In the opposite figure :

ABC is a triangle ,  $\overline{CD}$  bisects  $\angle C$  and intersects  $\overline{AB}$  at point D

$$, m(\angle BDC) = 100^\circ \text{ and } DB = DC$$

Prove that :

$$AC > DB$$




4

 In the opposite figure :

$\overrightarrow{AD} \parallel \overrightarrow{BC}$  ,  $m(\angle BAC) = 80^\circ$  and  $m(\angle DAC) = 30^\circ$

Prove that :

$BC > AB$




5

 In the opposite figure :

ABC is a triangle and  $D \in \overline{BC}$  where  $BD = AD$

Prove that :  $BC > AC$



**6 In the opposite figure :**


A is the midpoint of  $\overline{BD}$

$$, m(\angle ABC) = 70^\circ$$

$\therefore m(\angle ACB) = 50^\circ$

**Prove that :**

$$m(\angle D) > m(\angle DCA)$$





# Sheet (8)

## Triangle inequality

### Generally


In any triangle, the sum of the lengths of any two sides is greater than the length of the third side.

**i.e.** In any triangle such as  $\triangle ABC$

, we get :  $AB + BC > AC$

,  $BC + CA > AB$

,  $CA + AB > CB$



### Corollary

The length of any side in a triangle is greater than the difference between the lengths of the other two sides and less than their sum.


And you can prove that from the triangle inequality as follows :

In any triangle ABC :

$$AC + AB > BC \quad (1)$$

$$, \therefore AB + BC > AC \quad \text{i.e. } BC > AC - AB \quad (2)$$

From (1) and (2) , we deduce that :  $AC - AB < BC < AC + AB$



### Remark

To check the possibility that three lengths can be side lengths of a triangle, do as follows :

Compare the greatest length with the sum of the other two lengths :

- If the greatest length is greater than or equal to the sum of the other two lengths, you deduce that the three given lengths couldn't be lengths of the three sides of a triangle. (i.e. no triangle could be drawn with these side lengths).
- If the greatest length is less than the sum of the other two lengths, you deduce that the three given lengths could be lengths of the three sides of a triangle. (i.e. a triangle could be drawn with these side lengths).



**[1] Complete:**

- |           |   |
|-----------|---|
| <b>1</b>  | In any triangle the sum of any two sides ..... the length of the third side.  |
| <b>2</b>  | In $\Delta ABC$ , $AB + BC - AC > \dots\dots\dots$  |
| <b>3</b>  | The length of any side in a triangle ..... the sum of the lengths of the two other sides.   |
| <b>4</b>  | $ABC$ is a triangle , if $AB = 3$ cm. and $BC = 5$ cm. , then $AC \in ] \dots\dots\dots , \dots\dots\dots [$                                  |
| <b>5</b>  | $\Delta XYZ$ in which $XY = 4$ cm. and $YZ = 3$ cm. , then $XZ \in ] \dots\dots\dots , \dots\dots\dots [$                                     |
| <b>6</b>  | If $x$ cm. , 4 cm. and 5 cm. are lengths of the sides of a triangle , then $\dots\dots\dots < x < \dots\dots\dots$                            |
| <b>7</b>  | If : $x$ , 8 , 7 cm. are lengths of the sides of a triangle then $\dots\dots\dots < x < \dots\dots\dots$                                      |
| <b>8</b>  | If the lengths of two sides in triangle are 3 cm. and 9 cm. , then $\dots\dots\dots < \text{the length of third side} < \dots\dots\dots$      |
| <b>9</b>  | If the length of two sides of an isosceles triangle are 4 cm. and 10 cm. , then the length of the third side is $\dots\dots\dots$             |
| <b>10</b> | If the length of two sides in an isosceles triangle are 3 cm. and 7 cm. , then the length of the third side = $\dots\dots\dots$ cm.           |
| <b>11</b> | The length of two sides in an isosceles triangle are 8 cm. , 4 cm. then the length of the third side = $\dots\dots\dots$                      |
| <b>12</b> | If the lengths of two sides in an isosceles triangle are 6 cm. , 3 cm. , then the lengths of the third side is $\dots\dots\dots$ cm.          |
| <b>13</b> | The length of two sides in the isosceles triangle are 3 cm. and 8 cm. , then the length of third side equals $\dots\dots\dots$ cm.            |
| <b>14</b> | The triangle whose side lengths are $(2x - 1)$ cm. , $(x + 3)$ cm. , and 7 cm. becomes an equilateral triangle when $x = \dots\dots\dots$ cm. |

**[1] Choose the correct answer:**

- 1 The sum of lengths of any two sides in any triangle ..... the length of the third side.  
(a) is less than                      (b) is greater than                      (c) equals                      (d) otherwise
- 2 If the lengths of two sides in an isosceles triangle are 2 cm. and 5 cm. , then the length of the third side is ..... cm.  
(a) 2                      (b) 3                      (c) 5                      (d) 7
- 3  $\triangle ABC$  ,  $AB = 2$  cm. ,  $BC = 7$  cm. , then  $AC$  may equal .....  
(a) 2 cm.                      (b) 5 cm.                      (c) 9 cm.                      (d) 8 cm.
- 4 The numbers 6 , 3 , ..... can be lengths of sides of an isosceles triangle.  
(a) 3                      (b) 6                      (c) 9                      (d) 11
- 5 If the lengths of two sides in the isosceles triangle are 3 cm. , 7 cm. , then the length of third side = .....  
(a) 3 cm.                      (b) 7 cm.                      (c) 10 cm.                      (d) 4 cm.
- 6 If 3 cm. , 7 cm. are two side lengths in a triangle , then the smallest number of third side is .....  
(a) 3 cm.                      (b) 4 cm.                      (c) 5 cm.                      (d) 6 cm.
- 7 The numbers 5 , 4 and ..... can be lengths of sides of a triangle.  
(a) 8                      (b) 9                      (c) 10                      (d) 12
- 8 The numbers 4 , 8 , ..... can be lengths of sides of an isosceles triangle.  
(a) 4                      (b) 8                      (c) 12                      (d) 3
- 9 The lengths of two sides in a triangle are 4 cm. and 9 cm. and it has on axis of symmetry , then the length of third side is .....  
(a) 4 cm.                      (b) 5 cm.                      (c) 9 cm.                      (d) 13 cm.
- 10 The lengths of 5 cm. , 6 cm. and ..... can be length of the sides of a triangle.  
(a) 15 cm.                      (b) 13 cm.                      (c) 11 cm.                      (d) 8 cm.
- 11 The numbers 5 , 7 , ..... can be lengths of sides of triangle.  
(a) 12                      (b) 3                      (c) 2                      (d) 13



- 12** 3 , 10 , ..... can be lengths of sides of an isosceles triangle.  
(a) 10 (b) 8 (c) 6 (d) 4
- 13** In  $\triangle ABC$  if :  $AB = 3$  cm. and  $BC = 5$  cm. , then  $AC \in$  .....  
(a)  $]3 , 8]$  (b)  $[2 , 8]$  (c)  $]2 , 8 [$  (d)  $]2 , 5 [$
- 14** In the triangle ABC , if  $BC = 9$  cm. ,  $AB = 7$  cm. , then  $m(\angle C)$  .....  $m(\angle A)$   
(a) = (b)  $\geq$  (c)  $>$  (d)  $<$
- 15** Which of the following can be sides to draw the triangle .....  
(a) 5 cm. , 6 cm. , 12 cm. (b) 5 cm. , 6 cm. , 11 cm.  
(c) 5 cm. , 6 cm. , 4 cm. (d) 4 cm. , 6 cm. , 10 cm.
- 16** ) Which of the following numbers can be the lengths of sides of a triangle ?  
(a) 4 , 6 , 10 (b) 4 , 6 , 8 (c) 2 , 3 , 6 (d) 4 , 5 , 10
- 17** The lengths which can be the lengths of the sides of a triangle are .....  
(a) 3 , 4 , 7 (b) 3 , 3 , 6 (c) 3 , 5 , 7 (d) 1 , 5 , 7
- 18** Which of the following set of numbers can be lengths of sides of a triangle .....  
(a) 2 , 3 , 6 (b) 2 , 3 , 5 (c) 2 , 3 , 4 (d) 2 , 3 , 7
- 19** If the length of one side of a triangle is 5 cm. , then which of the following could be the lengths of the other two sides ?  
(a) 2 cm. and 3 cm. (b) 7 cm. and 2 cm.  
(c) 2 cm. and 2 cm. (d) 4 cm. and 6 cm.
- 20** Which of the following numbers cannot be the lengths of sides of a triangle .....  
(a) 7 , 7 , 5 (b) 9 , 9 , 9 (c) 3 , 6 , 12 (d) 3 , 4 , 5
- 21** In any triangle ABC :  $AB$  .....  $BC - AC$   
(a)  $>$  (b)  $<$  (c)  $=$  (d)  $\leq$
- 22** ) In the triangle ABC ,  $AC$  .....  $(AB - BC)$   
(a)  $>$  (b)  $\geq$  (c)  $\leq$  (d)  $<$

23 In any triangle ABC ,  $AB + BC \dots\dots\dots AC$

(a) =

(b) <

(c) >

(d)  $\leq$

## [2] Essay problems:


1

 In the opposite figure :


ABC is a triangle in which M is a point inside it.

Prove that :


$MA + MB + MC > \frac{1}{2}$  the perimeter of the triangle ABC



2

 Prove that the length of any side in a triangle is less than half of the perimeter.

3

 Prove that the sum of the lengths of two diagonals in a convex quadrilateral is less than its perimeter.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

